This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problems Mailbox.

Description Items Set TRACK? OR TRACE? OR TRACING OR MONITOR? 2941874 S1 (CLINICAL? OR MEDICAL) (5N) (STUDIES OR STUDY OR RESEARCH? OR 419717 \$2 TRIAL?) -5366 S1 (5N) S2 S3(15N) (SAMPLE? OR EXHIBIT? OR BLOOD? OR SERUM OR GENOTYP? S4 240 OR PHENOTYP?) S4 NOT PY>1999 100 S5 RD (unique items) 61 S6 ? show files 9:Business & Industry(R) Jul/1994-2003/Apr 24 File (c) 2003 Resp. DB Svcs. 15:ABI/Inform(R) 1971-2003/Apr 25 (c) 2003 ProQuest Info&Learning 16:Gale Group PROMT(R) 1990-2003/Apr 24 (c) 2003 The Gale Group File 148: Gale Group Trade & Industry DB 1976-2003/Apr 24 (c) 2003 The Gale Group File 160:Gale Group PROMT(R) 1972-1989 (c) 1999 The Gale Group File 275: Gale Group Computer DB (TM) 1983-2003/Apr 24 (c) 2003 The Gale Group File 621: Gale Group New Prod. Annou. (R) 1985-2003/Apr 24 (c) 2003 The Gale Group File 636:Gale Group Newsletter DB(TM) 1987-2003/Apr 24 (c) 2003 The Gale Group

(Item 1 from file: 9) 6/3, K/1

DIALOG(R)File 9:Business & Industry(R) (c) 2003 Resp. DB Svcs. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULLTEXT) 2605373 Supplier Number: 02605373 Diabetes Immunotherapy Clinical Trial Studies Nicotinamide for Diabetes (Italian study finds Nicotinamide (NA) can be beneficial to patients with recent-onset Type 1 diabetes)

World Disease Weekly Plus, p N/A

September 27, 1999

DOCUMENT TYPE: Newsletter (United States) LANGUAGE: English RECORD TYPE: Fulltext

WORD COUNT: 443

(USE FORMAT 7 OR 9 FOR FULLTEXT)

TEXT:

...carried out to optimize metabolic control as soon as possible after diagnosis and to maintain blood glucose level as near to normal as possible. Patient response to therapy was monitored throughout the study by investigating the occurrence of clinical (complete) remission defined, according to the recommendations of the International Diabetes Immunotherapy Group without any...

(Item 2 from file: 9) 6/3, K/2

DIALOG(R)File 9:Business & Industry(R) (c) 2003 Resp. DB Svcs. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULLTEXT) 2383152 Supplier Number: 02383152 Lovenox now available for acute deep-vein thrombosis treatment (FDA approved Lovenox as first low-molecular weight heparin for outpatient treatment of acute deep-vein thrombosis without pulmonary embolism)

Drug Store News, v 21, n 3, p CP10

February 15, 1999 DOCUMENT TYPE: Journal ISSN: 0191-7587 (United States)

LANGUAGE: English RECORD TYPE: Fulltext

WORD COUNT: 165

(USE FORMAT 7 OR 9 FOR FULLTEXT)

...given to the patient by continuous intravenous infusion, requiring several days of hospitalization and routine blood tests to monitor drug activity. According to a study done at Kaiser Permanente Medical Center, there was a clear economic advantage associated with using Lovenox in the outpatient setting...

(Item 3 from file: 9) 6/3.K/3

9:Business & Industry(R) DIALOG(R) File (c) 2003 Resp. DB Svcs. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULLTEXT) 2230884 Supplier Number: 02230884

Never richer

(An estimated 37 drug delivery products await approval; table lists upcoming drug delivery products with blockbuster potentials)

Med Ad News, v 17, n 8, p 38

August 1998

DOCUMENT TYPE: Journal ISSN: 0745-0907 (United States)

LANGUAGE: English RECORD TYPE: Fulltext

WORD COUNT: 3272

(USE FORMAT 7 OR 9 FOR FULLTEXT)

TEXT:

...generate at least \$100 million. Worn like a wristwatch, the GlucoWatch system is a painless, bloodless, and automatic glucose monitoring trials with GlucoWatch device. Cygnus moved into registration clinical at the end of 1997; the data from these trials are expected to...

(Item 4 from file: 9) 6/3, K/4

DIALOG(R)File 9:Business & Industry(R) (c) 2003 Resp. DB Svcs. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULLTEXT) 1229381 Supplier Number: 01229381

Chiron's pipeline

(Chiron Biocine's Gp120 anti-HIV vaccine is in Phase II clinical trials in

Med Ad News, v 14, n 7, p 20

July 1995

DOCUMENT TYPE: Journal ISSN: 0745-0907 (United States)

LANGUAGE: English RECORD TYPE: Fulltext

WORD COUNT: 616

(USE FORMAT 7 OR 9 FOR FULLTEXT)

TEXT:

...U.S. preclinical trials.

CMV bDNA Probe Test is being studied for quantifying viruses.

Critical blood analytes are in preclinical studies for monitoring critical care patients.

trials Discontinued clinical

Macrolin (macrophage colony stimulating factor) was in Phase III clinical trials in the United States...

(Item 1 from file: 15) 6/3, K/5

DIALOG(R)File 15:ABI/Inform(R)

(c) 2003 ProQuest Info&Learning. All rts. reserv.

01170603 98-19998

A community case management model

Faress, Mary Lynn

Nursing Management v27n3 PP: 65-66 Mar 1996

ISSN: 0744-6314 JRNL CODE: NSM

WORD COUNT: 651

... TEXT: admission compared to data before we implemented Care-Link Health status outcome measurements include (diabetic) blood glucose data, complication rates and co-morbidity tracking . Qualitative data encompass patient satisfaction studies .

Care-Link provides valuable expanded opportunities for BSN clinical students undergoing their community health rotation. The nursing faculty is currently considering...

(Item 2 from file: 15) 6/3, K/6

DIALOG(R)File 15:ABI/Inform(R)

(c) 2003 ProQuest Info&Learning. All rts. reserv.

00610156 92-25259

OSHA Targets AIDS in the Workplace

Kurland, Orin M.

Risk Management v39n4 PP: 16-18 Apr 1992

ISSN: 0035-5593 JRNL CODE: RMT

WORD COUNT: 1394

...TEXT: bundle of gauze and wrappers on the bed of an AIDS patient after taking a blood culture."

The February issue of the " Medical Monitor " reported Liability research suggesting that 88 percent of needlesticks are preventable. This finding simply further demonstrates the added...

(Item 1 from file: 16) 6/3, K/7

DIALOG(R)File 16:Gale Group PROMT(R)

(c) 2003 The Gale Group. All rts. reserv.

Supplier Number: 66680037 (USE FORMAT 7 FOR FULLTEXT) 08296274

Phase III Clinical Development.

BioPharm, pS18

Dec, 1999

Record Type: Fulltext Language: English

Document Type: Magazine/Journal; Trade

1503 Word Count:

subjects involved in a Phase III study depends on the disease being studied. Phase III trials are expensive: Clinical monitoring of patients can include X-rays, blood tests, and other laboratory studies in addition to physician fees and required hospitalizations. Whereas classical

(Item 2 from file: 16) 6/3, K/8

DIALOG(R)File 16:Gale Group PROMT(R)

(c) 2003 The Gale Group. All rts. reserv.

Supplier Number: 58033233 (USE FORMAT 7 FOR FULLTEXT)

New SND Solutions Framework Offers Business-Driven Applications To Extend the Functionality of Sales Automation Systems.

PR Newswire, p5422

Dec 6, 1999

Record Type: Fulltext Language: English

Document Type: Newswire; Trade

386 Word Count:

Solutions Framework

SND Solutions Framework delivers a complete platform of application modules, including Marketing Budget Tracking, Sample Allocation, Trials Management, and Literature Ordering, that can be enhanced to support client-specific business processes. The...

6/3,K/9 (Item 3 from file: 16)
DIALOG(R)File 16:Gale Group PROMT(R)
(c) 2003 The Gale Group. All rts. reserv.

06759895 Supplier Number: 55982143 (USE FORMAT 7 FOR FULLTEXT) Clinical Trial Studies Nicotinamide for Diabetes. (Statistical Data Included)

Marble, Michelle

World Disease Weekly Plus, pNA

Sept 27, 1999

Language: English Record Type: Fulltext

Article Type: Statistical Data Included

Document Type: Newsletter; Professional Trade

Word Count: 447

carried out to optimize metabolic control as soon as possible after diagnosis and to maintain **blood** glucose level as near to normal as possible. Patient response to therapy was **monitored** throughout the **study** by investigating the occurrence of **clinical** (complete) remission defined, according to the recommendations of the International Diabetes Immunotherapy Group without any...

6/3,K/10 (Item 4 from file: 16)
DIALOG(R)File 16:Gale Group PROMT(R)
(c) 2003 The Gale Group. All rts. reserv.

06682519 Supplier Number: 55918625 (USE FORMAT 7 FOR FULLTEXT) Micrologix initiates phase II clinical trials of MBI 226.

PR Newswire, p6263

Sept 30, 1999

Language: English Record Type: Fulltext

Document Type: Newswire; Trade

Word Count: 625

.. counter)

Micrologix is pleased to announce the commencement of a two-part US Phase II clinical trial as part of the fast track development program of MBI 226 for the prevention of central venous catheter-related bloodstream infection. The trial, developed in consultation with the US Food & Drug Administration (FDA), is designed...

6/3,K/11 (Item 5 from file: 16)
DIALOG(R)File 16:Gale Group PROMT(R)
(c) 2003 The Gale Group. All rts. reserv.

06454940 Supplier Number: 55082223 (USE FORMAT 7 FOR FULLTEXT) Firm's technology takes the work out of blood work. (VIA Medical Corp.)

Webb, Marion

San Diego Business Journal, v20, n25, p1(2)

June 21, 1999

Language: English Record Type: Fulltext Document Type: Magazine/Journal; Tabloid; Trade

Word Count: 1261

newborns, named VIA low-volume mode (LVM) neonatal systems.

Billman, who headed the 12-month clinical trial for the blood
gas monitoring system at Children's Hospital and holds a stake in the firm, touts the device...

(Item 6 from file: 16) 6/3,K/12 DIALOG(R) File 16: Gale Group PROMT(R) (c) 2003 The Gale Group. All rts. reserv.

Supplier Number: 55018249 (USE FORMAT 7 FOR FULLTEXT) Varian, Inc. Signs \$6 Million Sales Agreement With Seiko Instruments Inc. 'SII'; SII to Resell Analytical Instrument Line in Japan.

Business Wire, p0093

June 29, 1999

Language: English Record Type: Fulltext

Document Type: Newswire; Trade

Word Count: 417

for measuring the concentrations of a wide range of chemical elements in a variety of samples . Major applications include environmental monitoring, life science and clinical industrial quality assurance, geo-chemical exploration, and metallurgical analysis.

Varian, Inc.'s Vista system is...

(Item 7 from file: 16) 6/3, K/13DIALOG(R) File 16: Gale Group PROMT(R) (c) 2003 The Gale Group. All rts. reserv.

Supplier Number: 54592484 (USE FORMAT 7 FOR FULLTEXT) 06327555 Strategic Network Designs Launches Vertical Market Practice to Provide Sales Automation Solutions to the Pharmaceutical Industry.

Business Wire, p1106

May 10, 1999

Record Type: Fulltext Language: English

Document Type: Newswire; Trade

Word Count: 656

especially for the pharmaceutical industry. Working from a standard model, SND has developed solutions for Sample Ordering, Literature Trials Management and Marketing Budget Tracking Ordering, Clinical systems. This application model enables SND to provide clients with economies of scale in implementation...

(Item 8 from file: 16) 6/3,K/14 DIALOG(R) File 16: Gale Group PROMT(R)

(c) 2003 The Gale Group. All rts. reserv.

Supplier Number: 54073039 (USE FORMAT 7 FOR FULLTEXT) PE Biosystems Introduces System to Rapidly Identify Drug-Resistant HIV Strains.

Business Wire, p1056

March 11, 1999

Record Type: Fulltext Language: English

Document Type: Newswire; Trade

Word Count: 715

(USE FORMAT 7 FOR FULLTEXT)

TEXT:

...complete, automated system for standardized, sequence-based genotyping of resistance mutations in HIV. The HIV Genotyping System is expected to be of immediate use in investigational clinical trials to monitor the development of drug-resistant HIV strains and assess the effectiveness of anti-HIV drugs...

(Item 9 from file: 16) 6/3,K/15 DIALOG(R)File 16:Gale Group PROMT(R) (c) 2003 The Gale Group. All rts. reserv.

Supplier Number: 53723431 (USE FORMAT 7 FOR FULLTEXT) Enoxaparin cleared to treat DVT at home, in hospital. (deep-vein thrombosis)

Levy, Sandra

Drug Topics, v143, n2, p56(1)

Feb 1, 1999

Record Type: Fulltext Language: English

Document Type: Magazine/Journal; Trade

Word Count: 706

heparin, given by continuous intravenous infusion. This requires several days of hospitalization along with routine blood tests to monitor drug activity.

Two clinical trials comparing enoxaparin with unfractionated heparin found enoxaparin equivalent to adjusted-dose, continuous-infusion therapy with...

(Item 10 from file: 16) 6/3,K/16 DIALOG(R)File 16:Gale Group PROMT(R) (c) 2003 The Gale Group. All rts. reserv.

Supplier Number: 53099981 (USE FORMAT 7 FOR FULLTEXT) First Consumer Advertising Campaign for Prescription Weight-Management Product Launched.

PR Newswire, p4791

Oct 20, 1998

Record Type: Fulltext Language: English

Document Type: Newswire; Trade

1634 Word Count:

include dry mouth, headache, insomnia and constipation. Some people taking MERIDIA experienced substantial increases in blood pressure and heart rate. Regular blood pressure monitoring is required when prescribing MERIDIA. Clinical studies showed that these blood pressure and heart rate increases generally occur early on in treatment and the prescribing physician...

(Item 11 from file: 16) 6/3,K/17 DIALOG(R)File 16:Gale Group PROMT(R) (c) 2003 The Gale Group. All rts. reserv.

Supplier Number: 50256765 (USE FORMAT 7 FOR FULLTEXT) 05769071

EFFEXOR SAVES COSTS IN TREATING MAJOR DEPRESSION

Worldwide Biotech, v10, n9, pN/A

Sept 1, 1998

Record Type: Fulltext Language: English

Article Type: Article

Document Type: Newsletter; Trade

851 Word Count:

treated patients by the second week of treatment. Venlafaxine is associated with sustained increases in **blood** pressure in some patients, so regular **blood** -pressure **monitoring** is recommended. In **clinical trials** of Effexor XR, the overall incidence of sustained increases in **blood** pressure was 3%. Less than 1% of patients discontinued Effexor XR due to elevated blood...

6/3,K/18 (Item 12 from file: 16)
DIALOG(R)File 16:Gale Group PROMT(R)
(c) 2003 The Gale Group. All rts. reserv.

05711342 Supplier Number: 50172306 (USE FORMAT 7 FOR FULLTEXT)
Studies Show That the Antidepressant Effexor(R) XR Also Provides Effective
Relief of Anxiety Symptoms

PR Newswire, p715NYW019

July 15, 1998

Language: English Record Type: Fulltext

Article Type: Article

Document Type: Newswire; Trade

Word Count: 767

treated patients by the second week of treatment. Venlafaxine is associated with sustained increases in **blood** pressure in some patients, so regular **blood** -pressure **monitoring** is recommended. In **clinical trials** of Effexor XR, the overall incidence of sustained increases in **blood** pressure was 3%. Less than 1% of patients discontinued Effexor XR due to elevated blood...

6/3,K/19 (Item 13 from file: 16)
DIALOG(R)File 16:Gale Group PROMT(R)
(c) 2003 The Gale Group. All rts. reserv.

05706968 Supplier Number: 50164641 (USE FORMAT 7 FOR FULLTEXT)

Study Demonstrates That Effexor Saves Costs in Treating Major Depression.

Business Wire, p7131187

July 13, 1998

Language: English Record Type: Fulltext

Article Type: Article

Document Type: Newswire; Trade

Word Count: 888

treated patients by the second week of treatment. Venlafaxine is associated with sustained increases in **blood** pressure in some patients, so regular **blood** -pressure **monitoring** is recommended. In **clinical trials** of Effexor XR, the overall incidence of sustained increases in **blood** pressure was 3%. Less than 1% of patients discontinued Effexor XR due to elevated blood...

6/3,K/20 (Item 14 from file: 16)
DIALOG(R)File 16:Gale Group PROMT(R)
(c) 2003 The Gale Group. All rts. reserv.

04816408 Supplier Number: 47085524 (USE FORMAT 7 FOR FULLTEXT)
OMRON INTRODUCES WORLD'S SMALLEST BLOOD PRESSURE MONITOR
Biotech Equipment Update, v5, n2, pN/A

Feb 1, 1997

Language: English Record Type: Fulltext

Document Type: Newsletter; Trade

Word Count: 222

... numbers on the front of the unit.

Created by the market's leading manufacturer of blood pressure monitors and based on clinical research that has proven a direct relationship between blood pressure in the finger and the arm, the product is useful for those who need...

6/3,K/21 (Item 15 from file: 16)
DIALOG(R)File 16:Gale Group PROMT(R)
(c) 2003 The Gale Group. All rts. reserv.

04764619 Supplier Number: 47013900 (USE FORMAT 7 FOR FULLTEXT)
World's smallest blood pressure monitor; goes where you go, lets your
finger do the "talking".

Business Wire, p1061488

Jan 6, 1997

Language: English Record Type: Fulltext

Document Type: Newswire; Trade

Word Count: 249

... numbers on the front of the unit.

Created by the market's leading manufacturer of **blood** pressure **monitors** and based on **clinical research** that hasproven a direct relationship between **blood** pressure in the finger and the arm, the product is useful for those who need...

6/3,K/22 (Item 16 from file: 16)
DIALOG(R)File 16:Gale Group PROMT(R)
(c) 2003 The Gale Group. All rts. reserv.

04593827 Supplier Number: 46754396 (USE FORMAT 7 FOR FULLTEXT)

Have a Heart

Health Data Management, p62

Oct, 1996

Language: English Record Type: Fulltext

Document Type: Magazine/Journal; Trade

Word Count: 1483

... creates images of where the dyes are flowing through the body to identify problems in **blood** flow. The **clinical** system **monitoring** this **study** at a remote hospital creates live video images. New England's telecardiology system captures these...

6/3,K/23 (Item 17 from file: 16)
DIALOG(R)File 16:Gale Group PROMT(R)
(c) 2003 The Gale Group. All rts. reserv.

04078885 Supplier Number: 45940284

Biosensors are Making Steady Yet Limited Progress into the Marketplace

Genetic Engineering News, p14

Nov 15, 1995

Language: English Record Type: Abstract

Document Type: Magazine/Journal; Trade

ABSTRACT:

...with a sensor in order to detect the quantity of a specific compound in a sample . Biosensors are used in research , clinical testing, biprocess monitoring and environmental testing. Both Pharmacia and Yellow Springs Instrument Co. are offering new products. The...

(Item 18 from file: 16) 6/3, K/24DIALOG(R) File 16: Gale Group PROMT(R) (c) 2003 The Gale Group. All rts. reserv.

Supplier Number: 45825116 (USE FORMAT 7 FOR FULLTEXT) BIOPATH LAUNCHES LABELING AND EVENT TRACKING SYSTEM FOR CLINICAL TRIALS Biotech Equipment Update, v3, n10, pN/A

Oct 1, 1995

Record Type: Fulltext Language: English

Document Type: Newsletter; Trade

381 Word Count:

flexible. The product was designed specifically to meet the varying needs of pharmaceutical manufacturers during clinical trial protocols." BioPath's STICS (Sample Tracking and Information Control Systems) is a portfolio of PC-based labeling systems that combine standard...

(Item 19 from file: 16) DIALOG(R) File 16: Gale Group PROMT(R) (c) 2003 The Gale Group. All rts. reserv.

Supplier Number: 45277968 (USE FORMAT 7 FOR FULLTEXT) NORTHFIELD LABORATORIES REPORTS SECOND QUARTER RESULTS

PR Newswire, pN/A

Jan 20, 1995

Record Type: Fulltext Language: English

Document Type: Newswire; Trade

Word Count: 1232

of blood. There was no evidence of fever, no abnormal changes in heart rate or blood pressure, no vasoconstriction and no evidence of organ dysfunction in any recipient. trials continue to progress on track," said "Phase II clinical

Richard DeWoskin, Chairman and CEO. "With the success of our initial Phase II trials...

(Item 20 from file: 16) 6/3,K/26 DIALOG(R) File 16: Gale Group PROMT(R) (c) 2003 The Gale Group. All rts. reserv.

Supplier Number: 45084397 (USE FORMAT 7 FOR FULLTEXT)

Epigen plans private placement

Biotechnology Business News, pN/A

Oct 24, 1994

Language: English Record Type: Fulltext

Document Type: Newsletter; Trade

Word Count: 81

(USE FORMAT 7 FOR FULLTEXT)

TEXT:

... of its common stock in a private placement, and will use proceeds to

trials of a blood test for monitoring begin prospective clinical breast cancer patients. The company said it believes the private placement financing will prompt the...

(Item 21 from file: 16) 6/3,K/27 DIALOG(R) File 16:Gale Group PROMT(R) (c) 2003 The Gale Group. All rts. reserv.

Supplier Number: 44848818 (USE FORMAT 7 FOR FULLTEXT) 03472238 Treatment STATUS OF IMMUPATH REPORTED AS POTENTIAL AIDS TREATMENT AIDS Weekly, pN/A July 18, 1994 Record Type: Fulltext

Language: English

Document Type: Newsletter; General

Word Count: 487

3). (Decreasing CD4 cell concentrations are indicators of AIDS progression.)

The FDA has approved the study 's design, sample size, endpoints, clinical monitoring and manufacturing plan but has questioned the viral inactivation procedures used in processing Immupath. On...

(Item 22 from file: 16) 6/3,K/28 DIALOG(R) File 16: Gale Group PROMT(R) (c) 2003 The Gale Group. All rts. reserv.

Supplier Number: 44790045 (USE FORMAT 7 FOR FULLTEXT) Treatment News HEMACARE REPORTS ON REGULATORY STATUS OF IMMUPATH AIDS Weekly, pN/A June 27, 1994

Record Type: Fulltext Language: English

Document Type: Newsletter; General

Word Count: 473

CD4 lymphocyte concentrations of 50-200/cells per mm(3). The FDA has approved the study 's design, sample size, endpoints, monitoring and manufacturing plan but has questioned the viral clinical inactivation procedures used in processing Immupath. On...

(Item 23 from file: 16) 6/3,K/29 DIALOG(R) File 16: Gale Group PROMT(R) (c) 2003 The Gale Group. All rts. reserv.

Supplier Number: 44082083 (USE FORMAT 7 FOR FULLTEXT) 03007576 U.S. FDA IND GRANTED FOR IMMUNE RECONSTITUTION THERAPY AIDS Weekly, pN/A Sept 6, 1993 Record Type: Fulltext Language: English

Document Type: Newsletter; General

212 Word Count:

it with the FDA within six weeks. The expanded program will be administered as a monitored clinical trial ," Reiter concluded. CSI's immune reconstitution therapy includes an infusion of peripheral blood lymphocytes, which are specific types of white blood cells deficient in patients with AIDS. An...

(Item 24 from file: 16) 6/3,K/30 DIALOG(R) File 16: Gale Group PROMT(R) (c) 2003 The Gale Group. All rts. reserv.

Supplier Number: 43905281 02894016

Diabetes Report May Lift Sales of Medical Devices

The New York Times, pC4

June 15, 1993

Record Type: Abstract Language: English

Document Type: Newspaper; General

ABSTRACT:

Complications associated with diabetes could be prevented or delayed by more closely monitoring blood sugar levels, according to a new study . Medical technology companies that make monitoring equipment (Lifescan, for example) and insulin delivery systems (Minimed Technologies, for example) might benefit from...

(Item 25 from file: 16) 6/3, K/31DIALOG(R) File 16: Gale Group PROMT(R) (c) 2003 The Gale Group. All rts. reserv.

Supplier Number: 43676449 (USE FORMAT 7 FOR FULLTEXT) 02745736

Monitoring HIV Levels

Genesis Report-Dx, v2, n5, pN/A

March, 1993

Record Type: Fulltext Language: English

Document Type: Newsletter; Trade

Word Count: 342

journal Science.

Researchers at Genelabs Technologies believe the test could be used in therapeutic drug monitoring in clinical trials of AIDS treatments. By detecting the level of HIV in the blood , it could help gauge the effectiveness of antiviral agents in controlling AIDS. Genelabs is using...

(Item 26 from file: 16) 6/3,K/32 DIALOG(R) File 16: Gale Group PROMT(R) (c) 2003 The Gale Group. All rts. reserv.

Supplier Number: 43348785 (USE FORMAT 7 FOR FULLTEXT)

T Cell/Yamanouchi marketing deal Biotechnology Business News, pN/A

Oct 2, 1992

Record Type: Fulltext Language: English

Document Type: Newsletter; Trade

329 Word Count:

candidate, Trax CD4 Test Kit, in the near future. The product is already in late clinical trials in the US for monitoring white blood cells which express the protein receptor, CD4. The main target market for this will be...

(Item 27 from file: 16) 6/3,K/33 DIALOG(R) File 16: Gale Group PROMT(R) (c) 2003 The Gale Group. All rts. reserv. 02209489 Supplier Number: 42878160 (USE FORMAT 7 FOR FULLTEXT)

OSHA TARGETS AIDS IN THE WORKPLACE

Risk Management, p16

April, 1992

Language: English Record Type: Fulltext

Document Type: Magazine/Journal; Trade

Word Count: 1408

... bundle of gauze and wrappers on the bed of an AIDS patient after taking a blood culture."

The February issue of the "Medical Liability Monitor "reported research suggesting that 88 percent of needlesticks are preventable. This finding simply further demonstrates the added...

6/3,K/34 (Item 28 from file: 16)

DIALOG(R) File 16: Gale Group PROMT(R)

(c) 2003 The Gale Group. All rts. reserv.

02163769 Supplier Number: 42813312 (USE FORMAT 7 FOR FULLTEXT)

NEW PRODUCTS AND SERVICES --

Health Business, pN/A

March 6, 1992

Language: English Record Type: Fulltext

Document Type: Magazine/Journal; Trade

Word Count: 250

(USE FORMAT 7 FOR FULLTEXT)

TEXT:

...astigmatism. The holmium laser system currently is being evaluated to treat farsightedness -- A two-phase study finds that use of Wheaton Medical Technologies' Medi- Track electronic tracking system reduces blood pressure by enabling patients to follow medication dispensation schedules exactly. Patients using the pocket-sized...

6/3,K/35 (Item 29 from file: 16)

DIALOG(R) File 16: Gale Group PROMT(R)

(c) 2003 The Gale Group. All rts. reserv.

01831657 Supplier Number: 42315057 (USE FORMAT 7 FOR FULLTEXT)
THERMO ELECTRON TO ACQUIRE MANUFACTURER OF HEMOSTASIS MANAGEMENT PRODUCTS

PR Newswire, p1
August 26, 1991

Language: English Record Type: Fulltext Document Type: Newswire; Trade

Word Count: 230

PRODUCT NAMES: 3841280 (Surgical & Medical Blood Monitors);

8515200 (Applied Research)

6/3,K/36 (Item 30 from file: 16)

DIALOG(R) File 16: Gale Group PROMT(R)

(c) 2003 The Gale Group. All rts. reserv.

01532138 Supplier Number: 41868917 (USE FORMAT 7 FOR FULLTEXT)

Centocor's Day in the Sun

Health Business, v6, n7, pN/A

Feb 15, 1991

Record Type: Fulltext Language: English

Document Type: Magazine/Journal; Trade

287 Word Count:

of medical research and the flow of dollars from government and private sources. MRF features Medical Grant Monitor detailing every NIH research grant. To order or for a sample call 1-800/848-1153 or 202/828-4150 in Washington DC.

(Item 1 from file: 148) 6/3,K/37

DIALOG(R)File 148:Gale Group Trade & Industry DB (c) 2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 55807393 11358830

Colour doppler imaging for diagnosis of intracranial

hypotension. (Statistical Data Included)

Chao-Ching, Chen; Cheau-Lin, Luo; Shuu-Jiun, Wang; Chang-Ming, Chern;

Jong-Ling, Fuh; Sheng-Han, Lin; Han-Hwa, Hu

Lancet, 354, 9181, 826

Sept 4, 1999

DOCUMENT TYPE: Statistical Data Included ISSN: 0099-5355

RECORD TYPE: Fulltext LANGUAGE: English

LINE COUNT: 00314 WORD COUNT: 3397

Moreover, since changes in the blood flow in the superior ophthalmic vein were reversible after blood -patch treatment, CDFI would also be useful in monitoring the therapeutic effect of clinical management. However, this preliminary study included only a small number of participants, so further investigations in a larger population are...

(Item 2 from file: 148) 6/3,K/38

DIALOG(R) File 148: Gale Group Trade & Industry DB

(c)2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 55302919

Examining garlic's cardiovascular benefits.

Drug Store News, 21, 11, 53

July 19, 1999

RECORD TYPE: Fulltext LANGUAGE: English ISSN: 0191-7587

LINE COUNT: 00032 WORD COUNT: 362

aged garlic extract.

Included in the Wakunaga collection is a meta-analysis of some 16 clinical trials conducted between 1987 and 1994, tracking the effects of dry aged garlic extract on blood lipid levels. Sorting through the results of more than 900 hypercholesterolemic patients, according to the...

(Item 3 from file: 148) 6/3, K/39

DIALOG(R) File 148: Gale Group Trade & Industry DB

(c) 2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 53520315

What's new in hypertension: update on angiotensin II receptor

blockers. (includes related article on the role of angiotensin II in the development of hypertension)

Black, Henry R.

Consultant, 38, 11, 2715(7).

Nov, 1998

ISSN: 0010-7069 LANGUAGE: English RECORD TYPE: Fulltext; Abstract WORD COUNT: 3595 LINE COUNT: 00308

... receptor in vitro than losartan is, but it is unclear whether this feature has any clinical implications.(15)

EFFICACY

Clinical studies using ambulatory blood pressure monitoring have shown that all the ARBs produce smooth antihypertensive actions over a 24-hour period...

6/3,K/40 (Item 4 from file: 148)
DIALOG(R)File 148:Gale Group Trade & Industry DB
(c) 2003 The Gale Group. All rts. reserv.

10253927 SUPPLIER NUMBER: 20788184 (USE FORMAT 7 OR 9 FOR FULL TEXT)
MEASURING COMPLIANCE. (with medical treatment)

Fleming Jr., Harris

Drug Topics, v142, n11, p61(1)

June 1, 1998

ISSN: 0012-6616 LANGUAGE: English RECORD TYPE: Fulltext WORD COUNT: 620 LINE COUNT: 00055

... can work with them, and you can use them." Direct methods, used most commonly in **clinical research trials**, consist primarily of pharmacological **tracers** and **blood** and urine analysis. While they provide objective evidence, these methods can be very expensive and...

6/3,K/41 (Item 5 from file: 148)
DIALOG(R)File 148:Gale Group Trade & Industry DB
(c) 2003 The Gale Group. All rts. reserv.

09823347 SUPPLIER NUMBER: 19937356 (USE FORMAT 7 OR 9 FOR FULL TEXT)
Wyeth-Ayerst Receives U.S. Marketing Clearance For New, Once-Daily
Effexor(R) XR

PR Newswire, p1020NYM140

Oct 20, 1997

LANGUAGE: English RECORD TYPE: Fulltext WORD COUNT: 824 LINE COUNT: 00074

treated patients by the second week of treatment. Venlafaxine is associated with sustained increases in **blood** pressure in some patients, so regular **blood** -pressure **monitoring** is recommended. In **clinical** trials of Effexor XR, the overall incidence of sustained increases in **blood** pressure was 3%. Less than 1% of patients discontinued Effexor XR due to elevated blood...

6/3,K/42 (Item 6 from file: 148)
DIALOG(R)File 148:Gale Group Trade & Industry DB
(c)2003 The Gale Group. All rts. reserv.

09757179 SUPPLIER NUMBER: 19798382 (USE FORMAT 7 OR 9 FOR FULL TEXT)
Randomised trial of single and repeated fibrin glue compared with injection
of polidocanol in treatment of bleeding peptic ulcer.
Rutgeerts, Paul; Rauws, Erik; Wara, Pal; Swain, Paul; Hoos, Anton;
Solleder, Eva; Halttunen, Jorma; Dobrilla, Giorgio; Richter, Gerhard;

Prassler, Roland Lancet, v350, n9079, p692(5)

Sep 6, 1997

ISSN: 0099-5355 LANGUAGE: English RECORD TYPE: Fulltext; Abstract

WORD COUNT: 4260 LINE COUNT: 00356

... with the Declaration of Helsinki. Data were recorded carefully in case-record forms and the **study** was **monitored** according to good **clinical** practice regulations.33

Statistics The **sample** size of 250 people per treatment group was based on an expected recurrent bleeding rate...

6/3,K/43 (Item 7 from file: 148)
DIALOG(R)File 148:Gale Group Trade & Industry DB
(c) 2003 The Gale Group. All rts. reserv.

07616745 SUPPLIER NUMBER: 15949341 (USE FORMAT 7 OR 9 FOR FULL TEXT) Clinical trials with ambulatory blood pressure monitoring: fewer patients needed?

Staessen, Jan A.; Thijs, Lutgarde; Mancia, Giuseppe; Parati, Gianfranco; O'Brien, Eoin T.

Lancet, v344, n8936, p1552(5)

Dec 3, 1994

ISSN: 0099-5355 LANGUAGE: ENGLISH RECORD TYPE: FULLTEXT; ABSTRACT WORD COUNT: 3153 LINE COUNT: 00253

Clinical trials with ambulatory blood pressure monitoring: fewer patients needed?

... pressure reduction at a given time (eg, the time of peak plasma concentration) and the **blood** pressure changes at all other times.

It has also been suggested that ambulatory monitoring may facilitate the conduct of clinical trials by the early identification and exclusion of subjects whose blood pressure is raised only in the clinic environment (white coat reactors).[19,26] This feature...

6/3,K/44 (Item 8 from file: 148)
DIALOG(R)File 148:Gale Group Trade & Industry DB
(c)2003 The Gale Group. All rts. reserv.

07509433 SUPPLIER NUMBER: 15739728 (USE FORMAT 7 OR 9 FOR FULL TEXT) Pharmacological treatment of hypertension. (Hypertension Octet) Swales, J.D.

Lancet, v344, n8919, p380(6)

August 6, 1994

ISSN: 0099-5355 LANGUAGE: ENGLISH RECORD TYPE: FULLTEXT; ABSTRACT WORD COUNT: 5726 LINE COUNT: 00469

cardiac function in patients with hypertension. JAMA 1989; 261: 873-77. [24] Mancia G. Ambulatory blood pressure monitoring: research and clinical applications. J Hypertens 1990; 8 (suppl 7): s1-13. [25] Alderman MH. Blood pressure management: individualised treatment based on absolute risk and the potential for benefit. Ann Intern...

6/3,K/45 (Item 9 from file: 148)
DIALOG(R)File 148:Gale Group Trade & Industry DB
(c) 2003 The Gale Group. All rts. reserv.

07267830 SUPPLIER NUMBER: 15231319 (USE FORMAT 7 OR 9 FOR FULL TEXT) Hepatotoxic effects of tacrine administration in patients with Alzheimer's disease.

Watkins, Paul B.; Zimmerman, Hyman J.; Knapp, Margaret J.; Gracon, Stephen I.; Lewis, Karen W.

JAMA, The Journal of the American Medical Association, v271, n13, p992(7)

April 6, 1994

ISSN: 0098-7484 LANGUAGE: ENGLISH RECORD TYPE: FULLTEXT; ABSTRACT

WORD COUNT: 6434 LINE COUNT: 00522

TEXT:

observations, subsequent clinical studies have required frequent (weekly) monitoring of serum ALT levels.

6/3,K/46 (Item 10 from file: 148)

DIALOG(R) File 148: Gale Group Trade & Industry DB (c) 2003 The Gale Group. All rts. reserv.

06798843 SUPPLIER NUMBER: 14386838 (USE FORMAT 7 OR 9 FOR FULL TEXT)

Dexamethasone therapy for bacterial meningitis in children. (Swiss

Meningitis Study Group)

Schaad, Urs B.; Lips, Ueli; Gnehm, Hanspeter E.; Blumberg, Avihay; Heinzer, Ivo; Wedgwood, Joanna

Lancet, v342, n8869, p457(5)

August 21, 1993

ISSN: 0099-5355 LANGUAGE: ENGLISH RECORD TYPE: FULLTEXT; ABSTRACT WORD COUNT: 4166 LINE COUNT: 00345

... and at the end of antibiotic therapy.[17,18] Adverse or toxic reactions to the **study** drugs were sought by daily **clinical** observation and laboratory **monitoring** of complete **blood** cell count, electrolytes, creatinine, and liver enzymes in **serum**, glucose in urine, and occult blood in stool, on days 1,2, and 3, and...

6/3.K/47 (Item 11 from file: 148)

DIALOG(R) File 148: Gale Group Trade & Industry DB (c) 2003 The Gale Group. All rts. reserv.

06685537 SUPPLIER NUMBER: 13238305 (USE FORMAT 7 OR 9 FOR FULL TEXT)
CENTER FOR SPECIAL IMMUNOLOGY CLEARED TO PROVIDE IMMUNE RECONSTITUTION
THERAPY TO EXISTING PATIENTS; FDA APPROVES IND FOR LIMITED CLINICAL USE
PR Newswire, p0826LA013

August 26, 1993

LANGUAGE: ENGLISH RECORD TYPE: FULLTEXT WORD COUNT: 328 LINE COUNT: 00028

... it with the FDA within six weeks. The expanded program will be administered as a monitored clinical trial," Dr. Reiter concluded.

CSI's immune reconstitution therapy includes an infusion of peripheral **blood** lymphocytes, which are specific types of white blood cells deficient in patients with AIDS. An...

6/3,K/48 (Item 12 from file: 148)

DIALOG(R) File 148: Gale Group Trade & Industry DB (c) 2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 12562392

Long-term cardiac prognosis following noncardiac surgery.

Mangano, Dennis T.; Browner, Warren S.; Hollenberg, Milton; Li, Juliet; Tateo, Ida M.

JAMA, The Journal of the American Medical Association, v268, n2, p233(7)

July 8, 1992

RECORD TYPE: FULLTEXT; ABSTRACT LANGUAGE: ENGLISH ISSN: 0098-7484

LINE COUNT: 00470 WORD COUNT: 5498

minute.[3] Perioperatively, blood pressure was recorded from either an arterial line or an automated blood pressure cuff that was linked to a microcomputer. Heart rate was recorded from the Holter monitor .

Clinical Care

Clinical decisions were not controlled by study protocol. Routine clinical intraoperative monitoring included threelead ECG, pulse oximetry, and mass spectroscopy. Cardiac medications were continued to the morning...

(Item 13 from file: 148) 6/3,K/49 DIALOG(R)File 148:Gale Group Trade & Industry DB (c) 2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 12561500 Monitoring for myocardial ischemia during noncardiac surgery: a technology assessment of transesophageal echocardiography and 12-lead electrocardiography.

Eisenberg, Mark J.; London, Martin J.; Leung, Jacqueline M.; Browner, Warren S.; Hollenberg, Milton; Tubau, Julio F.; Tateo, Ida M.; Schiller, Nelson B.; Mangano, Dennis T. JAMA, The Journal of the American Medical Association, v268, n2, p210(7) July 8, 1992 RECORD TYPE: FULLTEXT; ABSTRACT LANGUAGE: ENGLISH ISSN: 0098-7484

LINE COUNT: 00594 6914 WORD COUNT:

continuous three-lead ECG, pulse oximetry, and mass spectroscopy were available to physicians. Intra-arterial blood pressure, central venous pressure, and pulmonaryartery pressure were also monitored if clinically indicated. Clinical decisions were unconstrained by study protocol.

Intraoperative Monitoring

All monitoring data were prospectively collected in a standard manner and were then analyzed after surgery (off...

(Item 14 from file: 148) DIALOG(R) File 148: Gale Group Trade & Industry DB (c)2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 12280406 Longitudinal evaluation of prostate-specific antigen levels in men with and without prostate disease.

Carter, H. Ballentine; Pearson, Jay D.; Metter, E. Jeffrey; Brant, Larry J. ; Chan, Daniel W.; Andres, Reubin; Fozard, James L.; Walsh, Patrick C. JAMA, The Journal of the American Medical Association, v267, n16, p2215(6)

April 22, 1992 RECORD TYPE: FULLTEXT; ABSTRACT LANGUAGE: ENGLISH ISSN: 0098-7484 LINE COUNT: 00365 WORD COUNT: 4556

case-control study may not be representative of changes that occur

in all men being monitored by PSA levels in clinical practice. In this study , PSA measurements were performed on stored serum samples and not on consecutively collected samples over many years as would be the case in clinical practice where the interassay variation...

(Item 15 from file: 148) 6/3,K/51

DIALOG(R) File 148: Gale Group Trade & Industry DB (c) 2003 The Gale Group. All rts. reserv.

SUPPLIER NUMBER: 12378268 (USE FORMAT 7 OR 9 FOR FULL TEXT) Ambulatory Monitoring and Blood Pressure Variability. (book reviews)

White, William B.

JAMA, The Journal of the American Medical Association, v267, n22, p3091(1)

June 10, 1992

DOCUMENT TYPE: review ISSN: 0098-7484 LANGUAGE: ENGLISH

RECORD TYPE: FULLTEXT

LINE COUNT: 00070 WORD COUNT: 839

with iiius, paper. \$99, ISBN 1-870026-82-9, London, England, Science Press. 1991.

Ambulatory **blood** pressure **monitoring** is a relatively new clinical and research technique that uses invasive and noninvasive blood pressure measurement methods combined with computer technology for data analysis to evaluate the variability of...

6/3, K/52(Item 16 from file: 148)

DIALOG(R) File 148: Gale Group Trade & Industry DB (c)2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 09301926 New uses and old for carbamazepine. (includes information on generics and on risk of birth defects)

Browne, Thomas R., III; Empting, Larry; Lydiard, R. Bruce; Yerby, Mark S. Patient Care, v24, n20, p48(15)

Dec 15, 1990

ISSN: 0031-305X LANGUAGE: ENGLISH

RECORD TYPE: FULLTEXT

WORD COUNT: 6629 LINE COUNT: 00562

that they are taking the drug). Dosage should be tapered when therapy is stopped.

The monitoring protocol used in many clinical trials of carbamazepine requires a complete blood count, liver function screen, and electrolyte determinations before starting treatment, then at the following intervals...

(Item 17 from file: 148) 6/3,K/53

DIALOG(R) File 148: Gale Group Trade & Industry DB

(c) 2003 The Gale Group. All rts. reserv.

SUPPLIER NUMBER: 09713935 (USE FORMAT 7 OR 9 FOR FULL TEXT) 04874357

FDA clears new heart imaging agent for emergency and routine use. (Cardiolite)

PR Newswire, pl227PH004

Dec 27, 1990

RECORD TYPE: FULLTEXT LANGUAGE: ENGLISH

LINE COUNT: 00080 935, WORD COUNT:

... to determine the efficacy of "Cardiolite" in detecting areas at risk of permanent damage. Worldwide clinical trials are under way to monitor its use in assessing the effectiveness of blood clot-dissolving therapies in coronary artery disease.

"Cardiolite" is available in more than 45 countries...

6/3,K/54 (Item 18 from file: 148)
DIALOG(R)File 148:Gale Group Trade & Industry DB
(c)2003 The Gale Group. All rts. reserv.

03940648 SUPPLIER NUMBER: 07038924 (USE FORMAT 7 OR 9 FOR FULL TEXT)

Average daily blood pressure, not office blood pressure, determines cardiac function in patients with hypertension.

White, William B.; Schulman, Peter; McCabe, Ellen J.; Dey, Holley M. JAMA, The Journal of the American Medical Association, v261, n6, p873(5) Feb 10, 1989

ISSN: 0098-7484 LANGUAGE: ENGLISH RECORD TYPE: FULLTEXT; ABSTRACT WORD COUNT: 4012 LINE COUNT: 00345

in the Hypertension Unit at the University of Connecticut Health Center, Farmington, who underwent ambulatory **blood** pressure **monitoring** for **clinical** or pharmacologic **research** purposes. Over a period of four years (1983 through 1987), 720 patients underwent ambulatory **blood** pressure monitoring and 77 of these patients were never previously treated with antihypertensive therapy. Twenty...

6/3,K/55 (Item 19 from file: 148)
DIALOG(R)File 148:Gale Group Trade & Industry DB
(c) 2003 The Gale Group. All rts. reserv.

03642701 SUPPLIER NUMBER: 07028837 (USE FORMAT 7 OR 9 FOR FULL TEXT)
Second National Conference on Alcohol Abuse and Alcoholism to be held in
San Diego Oct. 31 - Nov. 2. (NEWS ADVISORY)

PR Newswire, 1005DC010

Oct 5, 1988

LANGUAGE: ENGLISH RECORD TYPE: FULLTEXT WORD COUNT: 338 LINE COUNT: 00031

... conference boasts an expanded agenda: general sessions, five conference tracks, poster sessions, roundtables and an **exhibit** and resource center. The **tracks** are **research**; program administration; **clinical** issues; business and labor; and prevention, education and intervention.

Speakers from HHS will include Bowen...

6/3,K/56 (Item 1 from file: 160)
DIALOG(R)File 160:Gale Group PROMT(R)
(c) 1999 The Gale Group. All rts. reserv.

02314188

Clinical Data - Marketing Procedures
Annual Report 1989 p. 0

... heart have been referred to as "Silent Ischemia" and represent a transient reduced flow of **blood** to the cardiac tissue. With seventeen (17) years of experience in the **clinical** and **research** application of ambulatory ECG **monitoring**, the Company has been successful at developing

technology for detecting transient changes in the ambulatory...

6/3,K/57 (Item 1 from file: 621)

DIALOG(R) File 621: Gale Group New Prod. Annou. (R)

(c) 2003 The Gale Group. All rts. reserv.

01150650 Supplier Number: 41751309 (USE FORMAT 7 FOR FULLTEXT)

Du Pont Merck today reported that the Food and Drug Administration has cleared for use the first nonsurgical cardiac imaging agent that pinpoints heart attack damage, evaluates perfusion (blood flow) and determines pumping efficiency in a single study.

News Release, pN/A

Dec 27, 1990

Language: English Record Type: Fulltext

Document Type: Magazine/Journal; Trade

Word Count: 77.1

... to determine the efficacy of "Cardiolite" in detecting areas at risk of permanent damage. Worldwide clinical trials are under way to monitor its use in assessing the effectiveness of blood

clot-dissolving therapies in coronary artery disease.

"Cardiolite" is available in more than 45 countries...

6/3,K/58 (Item 1 from file: 636)

DIALOG(R)File 636:Gale Group Newsletter DB(TM)

(c) 2003 The Gale Group. All rts. reserv.

03955676 Supplier Number: 50311218 (USE FORMAT 7 FOR FULLTEXT)

And Almost Everything Else ...

Genesis Report-Dx, v7, n4, pN/A

Jan 1, 1998

Language: English Record Type: Fulltext

Document Type: Newsletter; Trade

Word Count: 6444

... without having a heavyweight marketing partner will probably be limited.

METABOLIC DISORDERS/DIABETES

Noninvasive glucose monitor from Cygnus performs well in clinical

Blood glucose monitoring: Cygnus, Inc (Redwood City, CA) has released favorable results from a series of research clinical...

6/3,K/59 (Item 2 from file: 636)

DIALOG(R) File 636: Gale Group Newsletter DB(TM)

(c) 2003 The Gale Group. All rts. reserv.

03766575 Supplier Number: 48153272 (USE FORMAT 7 FOR FULLTEXT)

And Also of Note ...

Genesis Report-Rx, v7, n1, pN/A

Dec 1, 1997

Language: English Record Type: Fulltext

Document Type: Newsletter; Trade

Word Count: 5896

and transient: dry mouth, headache, constipation, and insomnia. In some patients, however, Meridia substantially increases blood pressure, making regular monitoring necessary. In clinical trials, the number of patients who discontinued Meridia because of hypertension (0.4% for those on...

6/3,K/60 (Item 3 from file: 636)

DIALOG(R) File 636: Gale Group Newsletter DB(TM)

(c) 2003 The Gale Group. All rts. reserv.

03045365 Supplier Number: 46218388 (USE FORMAT 7 FOR FULLTEXT)

PRESENILIN GENE CONVICTED OF BOTH EARLY ONSET, LATE-ONSET ALZHEIMER'S

BIOWORLD Today, v7, n50, pN/A

March 12, 1996

Language: English Record Type: Fulltext

Document Type: Magazine/Journal; Trade

Word Count: 411

... Alzheimer's-type dementia, and 185 age-matched controls. In a separate cohort they took **blood** from 29 septuagenarian African-American patients of both sexes, plus controls.

All clinical centers contributing subjects to the **study** have a **track** record averaging 96-percent accuracy in diagnosing Alzheimer's, confirmed by post-mortem follow-ups...

6/3,K/61 (Item 4 from file: 636)

DIALOG(R) File 636: Gale Group Newsletter DB(TM) (c) 2003 The Gale Group. All rts. reserv.

01446057 Supplier Number: 41939750 (USE FORMAT 7 FOR FULLTEXT)

COFFEE INTAKE LINKED TO HEART ATTACK RISK IN STUDY

Food Chemical News, v33, n3, pN/A

March 18, 1991

Language: English Record Type: Fulltext

Document Type: Newsletter; Trade

Word Count: 447

... of the poster presentation. The risk was also independent of other factors such as age, serum cholesterol in medical school, and hypertension, the researchers said.

Tracking 1,040 former medical students, the Johns Hopkins researchers found relative coronary heart disease risks associated with five daily cups of coffee as 1...

```
Description
       Items
                TRACK? OR TRACE? OR TRACING OR MONITOR?
Set
                (CLINICAL OR MEDICAL) (5N) (STUDIES OR STUDY OR RESEARCH? OR
      2496039
S1
       381746
S2
             TRIAL?)
                S1(5N)S2
         4780
                S3(15N)(SAMPLE? OR EXHIBIT? OR BLOOD? OR SERUM OR GENOTYP?
S3
          229
S4
             OR PHENOTYP?)
                S4 NOT PY>1999
          101
S5
                RD (unique items)
           85
Sб
? show files
File 20:Dialog Global Reporter 1997-2003/Apr 25
         (c) 2003 The Dialog Corp.
File 476: Financial Times Fulltext 1982-2003/Apr 25
         (c) 2003 Financial Times Ltd
File 610:Business Wire 1999-2003/Apr 25
         (c) 2003 Business Wire.
File 613:PR Newswire 1999-2003/Apr 25
         (c) 2003 PR Newswire Association Inc
File 624:McGraw-Hill Publications 1985-2003/Apr 24
          (c) 2003 McGraw-Hill Co. Inc
File 634:San Jose Mercury Jun 1985-2003/Apr 24
          (c) 2003 San Jose Mercury News
File 810: Business Wire 1986-1999/Feb 28
          (c) 1999 Business Wire
File 813:PR Newswire 1987-1999/Apr 30
          (c) 1999 PR Newswire Association Inc
 File 442:AMA Journals 1982-2003/Aug B3
          (c)2003 Amer Med Assn -FARS/DARS apply
 File 149:TGG Health&Wellness DB(SM) 1976-2003/Apr W2
          (c) 2003 The Gale Group
 File 444: New England Journal of Med. 1985-2003/Apr W4
          (c) 2003 Mass. Med. Soc.
```

6/3,K/1 (Item 1 from file: 20)

DIALOG(R) File 20: Dialog Global Reporter (c) 2003 The Dialog Corp. All rts. reserv.

08576114 '(USE FORMAT 7 OR 9 FOR FULLTEXT)

New SND Solutions Framework Offers Business-Driven Applications To Extend the Functionality of Sales Automation Systems

PR NEWSWIRE

December 06, 1999

JOURNAL CODE: WPRW LANGUAGE: English RECORD TYPE: FULLTEXT

WORD COUNT: 415

(USE FORMAT 7 OR 9 FOR FULLTEXT)

... Solutions Framework

SND Solutions Framework delivers a complete platform of application modules, including Marketing Budget **Tracking**, **Sample**0 Allocation, Clinical **Trials** Management, and Literature Ordering, that can be enhanced to support client-specific business processes. The...

6/3,K/2 (Item 2 from file: 20)

DIALOG(R) File 20: Dialog Global Reporter

(c) 2003 The Dialog Corp. All rts. reserv.

07512894 (USE FORMAT 7 OR 9 FOR FULLTEXT)

Micrologix initiates phase II clinical trials of MBI 226

CANADA NEWSWIRE

September 30, 1999

JOURNAL CODE: WCNW LANGUAGE: English RECORD TYPE: FULLTEXT

WORD COUNT: 666

(USE FORMAT 7 OR 9 FOR FULLTEXT)

... counter)

Micrologix is pleased to announce the commencement of a two-part US Phase II clinical trial as part of the fast track development program of MBI 226 for the prevention of central venous catheter-related bloodstream infection. The trial, developed in consultation with the US Food & Drug Administration (FDA), is designed...

6/3,K/3 (Item 3 from file: 20)

DIALOG(R)File 20:Dialog Global Reporter

(c) 2003 The Dialog Corp. All rts. reserv.

05961799 (USE FORMAT 7 OR 9 FOR FULLTEXT)

Varian, Inc. Signs \$6 Million Sales Agreement With Seiko Instruments Inc. `SII'; SII to Resell Analytical Instrument Line in Japan

BUSINESS WIRE

June 29, 1999

JOURNAL CODE: WBWE LANGUAGE: English RECORD TYPE: FULLTEXT

WORD COUNT: 455

(USE FORMAT 7 OR 9 FOR FULLTEXT)

... for measuring the concentrations of a wide range of chemical elements in a variety of samples . Major applications include

Bode Akintola 25-Apr-03

environmental monitoring, life science and clinical studies, industrial quality assurance, geo-chemical exploration, and metallurgical analysis.

Varian, Inc.'s Vista system is...

6/3,K/4 (Item 4 from file: 20)

DIALOG(R) File 20: Dialog Global Reporter (c) 2003 The Dialog Corp. All rts. reserv.

05379111 (USE FORMAT 7 OR 9 FOR FULLTEXT)

Home Blood Pressure Telemonitoring Service Brings Reliable, Timely Data to
Doctors

PR NEWSWIRE

May 19, 1999

JOURNAL CODE: WPRW LANGUAGE: English RECORD TYPE: FULLTEXT

WORD COUNT: 710

(USE FORMAT 7 OR 9 FOR FULLTEXT)

... of readings, in a patient's normal environment. Dr. Pickering is presenting results of a **clinical trial** of home **blood** pressure **monitoring** with BPLink at the ASH meeting. Development and clinical testing of the BPLink system has...

6/3,K/5 (Item 5 from file: 20)

DIALOG(R)File 20:Dialog Global Reporter (c) 2003 The Dialog Corp. All rts. reserv.

05252907 (USE FORMAT 7 OR 9 FOR FULLTEXT)

Strategic Network Designs Launches Vertical Market Practice to Provide Sales Automation Solutions to the Pharmaceutical Industry

BUSINESS WIRE

May 10, 1999

JOURNAL CODE: WBWE LANGUAGE: English RECORD TYPE: FULLTEXT

WORD COUNT: 850

(USE FORMAT 7 OR 9 FOR FULLTEXT)

especially for the pharmaceutical industry. Working from a standard model, SND has developed solutions for **Sample** Ordering, Literature Ordering, **Clinical Trials** Management and Marketing Budget **Tracking** systems. This application model enables SND to provide clients with economies of scale in implementation...

6/3,K/6 (Item 6 from file: 20)

DIALOG(R) File 20: Dialog Global Reporter

(c) 2003 The Dialog Corp. All rts. reserv.

04604093 (USE FORMAT 7 OR 9 FOR FULLTEXT)

PE Biosystems Introduces System to Rapidly Identify Drug-Resistant HIV Strains

BUSINESS WIRE

March 11, 1999

JOURNAL CODE: WBWE LANGUAGE: English RECORD TYPE: FULLTEXT

WORD COUNT: 865

... complete, automated system for standardized, sequence-based

genotyping of resistance mutations in HIV. The HIV Genotyping System is expected to be of immediate use in investigational clinical trials to monitor the development of drug-resistant HIV strains and assess the effectiveness of anti-HIV drugs...

6/3,K/7 (Item 7 from file: 20)

DIALOG(R) File 20: Dialog Global Reporter (c) 2003 The Dialog Corp. All rts. reserv.

03167963 (USE FORMAT 7 OR 9 FOR FULLTEXT)

First Consumer Advertising Campaign for Prescription Weight-Management Product Launched

PR NEWSWIRE

October 20, 1998

JOURNAL CODE: WPRW LANGUAGE: English RECORD TYPE: FULLTEXT

WORD COUNT: 1448

(USE FORMAT 7 OR 9 FOR FULLTEXT)

include dry mouth, headache, insomnia and constipation. Some people taking MERIDIA experienced substantial increases in **blood** pressure and heart rate. Regular **blood** pressure **monitoring** is required when prescribing MERIDIA. **Clinical studies** showed that these **blood** pressure and heart rate increases generally occur early on in treatment and the prescribing physician...

6/3,K/8 (Item 8 from file: 20)

DIALOG(R)File 20:Dialog Global Reporter (c) 2003 The Dialog Corp. All rts. reserv.

02343974 (USE FORMAT 7 OR 9 FOR FULLTEXT)

Health Canada Approves Treatment For Suppression Of Genital Herpes

CANADA NEWSWIRE

July 28, 1998 9:51

JOURNAL CODE: WCNW LANGUAGE: English RECORD TYPE: FULLTEXT

WORD COUNT: 889

(USE FORMAT 7 OR 9 FOR FULLTEXT)

... said Dr. Diaz-Mitoma. ``Twice-daily dosing allows the drug to continually circulate in the **bloodstream** so there is a greater potential to stop the virus in its **tracks**,'' he continued.

One-Year Clinical Trial

In a subsequent one-year, multi-centre study, Famvir(R) 250 mg taken twice-daily...

6/3,K/9 (Item 9 from file: 20)

DIALOG(R) File 20: Dialog Global Reporter

(c) 2003 The Dialog Corp. All rts. reserv.

02215247 (USE FORMAT 7 OR 9 FOR FULLTEXT)

Studies Show That the Antidepressant Effexor(R) XR Also Provides Effective Relief of Anxiety Symptoms

PR NEWSWIRE

July 15, 1998 9:21

JOURNAL CODE: WPRW LANGUAGE: English RECORD TYPE: FULLTEXT

WORD COUNT: 747

(USE FORMAT 7 OR 9 FOR FULLTEXT)

treated patients by the second week of treatment. Venlafaxine is associated with sustained increases in **blood** pressure in some patients, so regular blood -pressure monitoring is recommended. In clinical trials of Effexor XR, the overall incidence of sustained increases in pressure was 3%. Less than 1% of patients discontinued Effexor XR blood due to elevated blood...

(Item 10 from file: 20) 6/3, K/10DIALOG(R)File 20:Dialog Global Reporter (c) 2003 The Dialog Corp. All rts. reserv.

02187793 (USE FORMAT 7 OR.9 FOR FULLTEXT) Study Demonstrates That Effexor Saves Costs in Treating Major Depression BUSINESS WIRE LANGUAGE: English RECORD TYPE: FULLTEXT 9:26 July 13, 1998 JOURNAL CODE: WBWE WORD COUNT: 873

(USE FORMAT 7 OR 9 FOR FULLTEXT)

treated patients by the second week of treatment. Venlafaxine is associated with sustained increases in blood pressure in some patients, so regular blood -pressure monitoring is recommended. In clinical of Effexor XR, the overall incidence of sustained increases in blood pressure was 3%. Less than 1% of patients discontinued Effexor XR due to elevated blood...

(Item 11 from file: 20) 6/3,K/11 DIALOG(R)File 20:Dialog Global Reporter (c) 2003 The Dialog Corp. All rts. reserv.

01809632 (USE FORMAT 7 OR 9 FOR FULLTEXT)

Show Effexor(R) XR Produces Significantly Higher Depression Studies Remission Rates Than Prozac

PR NEWSWIRE

JOURNAL CODE: WPRW LANGUAGE: English RECORD TYPE: FULLTEXT

WORD COUNT: 659

(USE FORMAT 7 OR 9 FOR FULLTEXT)

placebo-treated patients by the second weekof treatment. Venlafaxine is associated with sustained increases in bloodpressure in some patients, so regular 1blood -pressure monitoring is recommended. In of Effexor XR, the overall incidence of sustained blood pressure was 3%. Less than 1% of patients discontinued clinical increasesin Effexor XR dueto elevated blood pressure...

(Item 1 from file: 476) DIALOG(R) File 476: Financial Times Fulltext 6/3,K/12 (c) 2003 Financial Times Ltd. All rts. reserv.

0008505605 BOGFEAAADYFT

News: UK: Watchdog attacks NHS training

MARK SUZMAN, Social Affairs Correspondent Financial Times, London Edition 1 ED, P 12

Wednesday, June 5, 1996

DOCUMENT TYPE: Stories; NEWSPAPER LANGUAGE: ENGLISH RECORD TYPE:

FULLTEXT

Word Count: 341

...government to ensure that training for and monitoring of clinical activities is properly implemented and monitored . The previous study had been criticised by medical groups as unrepresentative due to its small sample size.

According to Dr Jonathan Boyce, the director of health studies at the commission, the...

(Item 1 from file: 634) 6/3, K/13

DIALOG(R) File 634: San Jose Mercury

(c) 2003 San Jose Mercury News. All rts. reserv.

06177047

DEVICE COULD SAVE DIABETICS FROM BLOODLETTING

San Jose Mercury News (SJ) - Tuesday June 25, 1991

By: Associated Press

Edition: Morning Final Section: Science & Medicine Page: 2C

Word Count: 196

...research director of Futrex Inc. in Gaithersburg, Md.

"If the technology is proven effective in clinical trials , blood monitoring could be pain-free and far more convenient," alucose Rosenthal said.

An estimated 14 million Americans...

(Item 2 from file: 634) 6/3, K/14

DIALOG(R)File 634:San Jose Mercury

(c) 2003 San Jose Mercury News. All rts. reserv.

03579159

TWO OF A KIND -- RELATIVELY SPEAKING RESEARCH ON TWINS FINDS ASTONISHING 'COINCIDENCES' OF NAMES, BEHAVIOR, JOBS

SAN JOSE MERCURY NEWS (SJ) - Thursday, October 2, 1986

By: PAULA FROKE, Associated Press

Edition: Stock Final Section: Front Page: 2A

Word Count: 998

...other studies.''

During their week at the university, the twins are asked to provide medical histories. Researchers monitor their hearts, blood complete blood pressure, allergies, and motor skills such as reflexes, hand steadiness and hand-eye coordination.

Other...

(Item 1 from file: 810) 6/3,K/15 DIALOG(R) File 810: Business Wire

Bode Akintola 25-Apr-03

(c) 1999 Business Wire . All rts. reserv.

0658870 BW1488

OMRON HEALTHCARE 2: World's smallest blood pressure monitor; goes where you go, lets your finger do the "talking"

January 06, 1997

Business Editors, Health & Medical Writers Byline:

...numbers on the front of the unit.

Created by the market's leading manufacturer of **blood** pressure monitors and based on clinical research that has proven a direct relationship between blood pressure in the finger and the arm, the product is useful for those who need...

(Item 2 from file: 810) 6/3,K/16

DIALOG(R) File 810: Business Wire

(c) 1999 Business Wire . All rts. reserv.

0412467 BW035

HemaCare reports on regulatory status of Immupath --HEMACARE CORP: potential AIDS treatment

June 16, 1994

Health/Medical Writers Byline:

(Decreasing CD4 cell concentrations are

indicators of AIDS progression.)

The FDA has approved the study 's design, sample size, endpoints, monitoring and manufacturing plan but has questioned the viral clinical

inactivation procedures used in processing Immupath. On...

(Item 3 from file: 810) 6/3,K/17

DIALOG(R) File 810: Business Wire

(c) 1999 Business Wire . All rts. reserv.

BW045 0378414

TheraTech completes phase III clinical trials of testosterone patch for the treatment of male hormone deficiency

January 12, 1994

Medical/Business Editors Byline:

testosterone patch for periods lasting six to 12 months, and were evaluated by measuring serum hormone levels, assessing hypogonadal symptoms and monitoring safety parameters.

The studies were conducted at medical centers in the United States and Sweden, including the Johns Hopkins School of Medicine, the...

6/3,K/18 (Item 4 from file: 810)
DIALOG(R)File 810:Business Wire

(c) 1999 Business Wire . All rts. reserv.

0339096 BW729

CHROMATICS CLR SCI INTL: Investors Associates initiates coverage of Chromatics

June 10, 1993

Byline:

Business Editors

...doctor would merely scan the infant's skin and receive a diagnosis that has, in **clinical research studies**, **tracked** in direct ratio to **blood** testing. The company, working in conjunction with Mt. Sinai Hospital, began the trials in 1988...

6/3,K/19 (Item 5 from file: 810)
DIALOG(R)File 810:Business Wire
(c) 1999 Business Wire . All rts. reserv.

0199934 BW728

CLINICAL DATA: Clinical Data sells EnviroMed subsidiary to Browning Ferris Industries

October 31, 1990

Byline: Business Editors

...subsidiary, Vital
Scientific NV, will continue to manufacture photometric instrumentation used in the analysis of blood chemistry. The Drug Research Service Division will continue to provide Ambulatory ECG and Blood Pressure monitoring services to pharmaceutical companies and medical research institutes in the United States and Europe.

Israel M. Stein, M.D., president of Clinical...

6/3,K/20 (Item 6 from file: 810)
DIALOG(R) File 810: Business Wire
(c) 1999 Business Wire . All rts. reserv.

0162583 BW745

CLINICAL DATA 2: Clinical Data secures stockholder approval and closes divestiture

February 7, 1990

Byline: Business Editors/Medical Writers

...Data, through its Drug Research Services Division will continue to provide Ambulatory ECG (Holter) and **Blood** Pressure monitoring services to pharmaceutical companies and medical research institutes. The Drug Research Services Division works with 15 international drug companies, performs studies in the United States,

Canada...

6/3,K/21 (Item 7 from file: 810)

DIALOG(R) File 810: Business Wire

(c) 1999 Business Wire . All rts. reserv.

0159291 BW713

CLINICAL DATA: Clinical Data and United Medical sign definitive agreement

January 16, 1990

Byline: Business Editors

...Drug Research Services Division of Clinical
Data will continue to provide Ambulatory ECG (Holter) and Blood
Pressure monitoring services to pharmaceutical companies and medical
research institutes in the United States and Europe. Clinical Data
will also, through its Dutch subsidiary...

6/3,K/22 (Item 8 from file: 810)

DIALOG(R) File 810: Business Wire

(c) 1999 Business Wire . All rts. reserv.

0148314 BW901

UNITED MEDICAL: United Medical signs letter to acquire Clinical Data Inc.

October 19, 1989

Byline: Business Editors And Medical Writers

...operation, its research division provides complete
data analysis and management services of Holter and ambulatory blood
pressure monitoring studies for pharmaceutical companies worldwide.
United Medical Corp.'s other health-care services include physical

United Medical Corp.'s other health-care services include physical examinations, electronic new drug applications for submission...

6/3,K/23 (Item 9 from file: 810)

DIALOG(R) File 810: Business Wire

(c) 1999 Business Wire . All rts. reserv.

0148300 BW880

CLINICAL DATA: (UM) Clinical Data's Diagnostic Services Division to be acquired by United Medical

October 19, 1989

Byline: Business Editors

...the Drug Research Services Division of Clinical
Data will continue to provide Ambulatory ECG and Blood Pressure
monitoring services to pharmaceutical companies and medical research
institutes in the United States and Europe.

Clinical Data will through its Dutch subsidiary, Vital...

6/3, K/24 (Item 1 from file: 813)

DIALOG(R) File 813: PR Newswire

(c) 1999 PR Newswire Association Inc. All rts. reserv.

1170689

NYM140

Wyeth-Ayerst Receives U.S. Marketing Clearance For New, Once-Daily Effexor(R) XR

DATE: October 20, 1997 15:25 EDT WORD COUNT: 738

... treated patients by the second week of treatment. Venlafaxine is associated with sustained increases in **blood** pressure in some patients, so regular **blood** -pressure **monitoring** is recommended. In **clinical trials** of Effexor XR, the overall incidence of sustained increases in **blood** pressure was 3%. Less than 1% of patients discontinued Effexor XR due to elevated blood...

6/3,K/25 (Item 2 from file: 813)

DIALOG(R) File 813: PR Newswire

(c) 1999 PR Newswire Association Inc. All rts. reserv.

0780508

CL011

NORTHFIELD LABORATORIES REPORTS SECOND QUARTER RESULTS

DATE: January 20, 1995 10:55 EST WORD COUNT: 1,121

...of blood. There was no evidence of fever, no abnormal changes in heart rate or **blood** pressure, no vasoconstriction and no evidence of organ dysfunction in any recipient.

"Phase II clinical trials continue to progress on track," said Richard DeWoskin, Chairman and CEO. "With the success of our initial Phase II trials...

6/3,K/26 (Item 3 from file: 813)

DIALOG(R) File 813:PR Newswire

(c) 1999 PR Newswire Association Inc. All rts. reserv.

0625930 LA013

CENTER FOR SPECIAL IMMUNOLOGY CLEARED TO PROVIDE IMMUNE RECONSTITUTION THERAPY TO EXISTING PATIENTS; FDA APPROVES IND FOR LIMITED CLINICAL USE

DATE: August 26, 1993 09:58 EDT WORD COUNT: 286

...it with the FDA within

six weeks. The expanded program will be administered as a monitored clinical trial," Dr. Reiter concluded.

CSI's immune reconstitution therapy includes an infusion of peripheral **blood** lymphocytes, which are specific types of white blood cells deficient in patients with AIDS. An...

6/3,K/27 (Item 4 from file: 813)

DIALOG(R)File 813:PR Newswire

(c) 1999 PR Newswire Association Inc. All rts. reserv.

0332076 PH004

FDA CLEARS NEW HEART IMAGING AGENT FOR EMERGENCY AND ROUTINE USE

DATE: December 27, 1990 09:29 EST WORD COUNT: 854

...to determine the efficacy of "Cardiolite" in detecting areas at risk of permanent damage. Worldwide clinical trials are under way to monitor its use in assessing the effectiveness of blood clot-dissolving therapies in coronary artery disease.

"Cardiolite" is available in more than 45 countries...

6/3,K/28 (Item 5 from file: 813)

DIALOG(R) File 813: PR Newswire

(c) 1999 PR Newswire Association Inc. All rts. reserv.

0109942 DC010A

SECOND NATIONAL CONFERENCE ON ALCOHOL ABUSE AND ALCOHOLISM TO BE HELD IN SAN DIEGO OCT. 31 - NOV. 2

DATE: October 5, 1988 12:12 E.T. WORD COUNT: 324

...conference boasts an expanded agenda: general sessions, five conference tracks, poster sessions, roundtables and an **exhibit** and resource center. The **tracks** are **research**; program administration; **clinical** issues; business and labor; and prevention, education and intervention.

Speakers from HHS will include Bowen...

6/3,K/29 (Item 1 from file: 442)

DIALOG(R) File 442:AMA Journals (c) 2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00108858

COPYRIGHT American Medical Association 1998

Clinical Research Designs for Emerging Treatments for Alzheimer Disease Moving Beyond Placebo-Controlled Trials (ARTICLE)

KNOPMAN, DAVID; KAHN, JEFFREY; MILES, STEVEN

Archives of Neurology

Nov, 1998; Special: tzn1425

LINE COUNT: 00473

... results.11/ Tacrine therapy requires 4 times a day dosing and either weekly or biweekly blood test monitoring (for transaminase elevations). In clinical trials , more than 50% of patients receiving high-dose tacrine stopped taking it before the end...

6/3,K/30 (Item 2 from file: 442)

DIALOG(R) File 442: AMA Journals

(c) 2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00106394

COPYRIGHT American Medical Association 1998

Coronary Artery Disease in Women A Historical Perspective (ARTICLE)

THOMAS, JOAN L.; BRAUS, PATRICIA A.

Archives of Internal Medicine

Feb 23,, 1998; Review Article: tzi333

LINE COUNT: 00484

... million to establish the Office of Research on Women's Health at the NIH, to monitor the inclusion of women in clinical trials funded by

A second National Heart, Lung, and Blood Institute conference the agency.6,24/ examining women and heart disease took place in 1992.23/ Led by Dr...

(Item 3 from file: 442) 6/3,K/31

DIALOG(R) File 442: AMA Journals

(c)2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

COPYRIGHT American Medical Association 1995

Is Methylphenidate Like Cocaine? Studies on Their Pharmacokinetics and Distribution in the Human Brain (ARTICLE) .

VOLKOW, NORA D.; DING, YU-SHIN; FOWLER, JOANNA S.; WANG, GENE-JACK; LOGAN, JEAN; GATLEY, JOHN S.; DEWEY, STEPHEN; ASHBY, CHARLES; LIEBERMANN, JEFFREY; HITZEMANN, ROBERT; WOLF, ALFRED P.

Archives of General Psychiatry

June, 1995; Original Article: ps_456

LINE COUNT: 00676

... throughout the study, and selected plasma samples were analyzed for the presence of unchanged radiotracer. Blood sampling was performed in the same manner as that for the human studies. Vitalsigns were monitored throughout the length of the study (90303B monitor, SpaceLabs Medical , Redmond, Wash). Details about the procedures have been published./23/ DATA ANALYSES

Regions of interest...

(Item 4 from file: 442) 6/3,K/32

DIALOG(R)File 442:AMA Journals

(c)2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00053995

COPYRIGHT American Medical Association 1992

Long-term cardiac prognosis following noncardiac surgery.

Mangano, Dennis T.; Browner, Warren S.; Hollenberg, Milton; Li, Juliet; Tateo, Ida M.

JAMA, The Journal of the American Medical Association

July 8, 1992; v268: p233(7)

LINE COUNT: 00481

... minute./3/ Perioperatively, blood pressure was recorded from either an arterial line or an automated **blood** pressure cuff that was linked to a microcomputer. Heart rate was recorded from the Holter monitor .

Clinical decisions were not controlled by study protocol. Routine Clinical Care

intraoperative monitoring included threelead ECG, pulse clinical oximetry, and mass spectroscopy. Cardiac medications were continued to the morning...

(Item 5 from file: 442) 6/3,K/33

DIALOG(R) File 442: AMA Journals

(c)2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00053991

COPYRIGHT American Medical Association 1992

Monitoring for myocardial ischemia during noncardiac surgery: a technology transesophageal echocardiography and 12-lead of assessment electrocardiography.

Eisenberg, Mark J.; London, Martin J.; Leung, Jacqueline M.; Browner, Warren S.; Hollenberg, Milton; Tubau, Julio F.; Tateo, Ida M.; Schiller,

Nelson B.; Mangano, Dennis T.

JAMA, The Journal of the American Medical Association

July 8, 1992; v268: p210(7)

LINE COUNT: 00605

... continuous three-lead ECG, pulse oximetry, and mass spectroscopy were available to physicians. Intra-arterial blood pressure, central venous pressure, and pulmonaryartery pressure were also monitored if clinically indicated. Clinical decisions were unconstrained by study protocol.

Intraoperative Monitoring

data were prospectively collected in a standard monitoring manner and were then analyzed after surgery (off...

(Item 6 from file: 442) 6/3,K/34

DIALOG(R) File 442:AMA Journals

(c)2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00053779

COPYRIGHT American Medical Association 1992

Ambulatory Monitoring and Blood Pressure Variability._(book reviews) (review)

White, William B.

JAMA, The Journal of the American Medical Association

June 10, 1992; v267: p3091(1)

LINE COUNT: 00074

... with iiius, paper. \$99, ISBN 1-870026-82-9, London, England, Science Press. 1991.

pressure monitoring is a relatively new Ambulatory **blood** technique that uses invasive and noninvasive clinical and research blood pressure measurement methods combined with computer technology for data analysis to evaluate the variability of...

(Item 7 from file: 442) 6/3,K/35

DIALOG(R) File 442: AMA Journals

(c) 2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00052696

The Impact of Long-term Warfarin Therapy on Quality of Life: Evidence From a Randomized Trial (Article)

Lancaster, Tim R., MRCP; Singer, Daniel E., MD; Sheehan, Mary A., RN; Oertel, Lynn B., RN-C, MS; Maraventano, Sue W., RN; Hughes, Robert A., MD; Kistler, J. Philip, MD Archives of Internal Medicine 1991; 151: 1944 (6)

...of the increased risk of bleeding conferred by warfarin and also because of the frequent **blood** tests and close medical follow-up needed to monitor warfarin therapy. Randomized clinical trials are providing valid estimates of warfarin's efficacy and risk of hemorrhage. Our work assessed...

6/3,K/36 (Item 8 from file: 442)
DIALOG(R)File 442:AMA Journals
(c)2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00052683

Growth Hormone Therapy in Hypophosphatemic Rickets (Article)

Wilson, Darrel M., MD; Lee, Phillip D. K., MD; Morris, Alan H., MD; Reiter, Edward O., MD; Gertner, Joseph M., MB, MCRP; Marcus, Robert, MD; Quarmby, Valerie E., PhD; Rosenfeld, Ron G., MD American Journal of Diseases of Children 1991; 145: 1165 (6)

... of the study. Nutritional intake of calcium, phosphorus, protein, carbohydrate, fat, and total energy was **monitored** by the **Clinical Research** Center dietitian at each institution. Assays Growth hormone levels were determined by immunoassay. **Serum** and urine phosphate and calcium levels, serum alkaline phosphatase levels, and other biochemical measures were...

6/3,K/37 (Item 9 from file: 442)
DIALOG(R)File 442:AMA Journals
(c)2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00043313 Copyright (C) 1988 American Medical Association

Pepsin Release by Prostaglandin El Analogue; A Potential Therapeutic Problem

BASSON, MARC D.; ZUCKER, KARL A.; ADRIAN, THOMAS E.; ZDON, MICHAEL J.; BALLANTYNE, GARTH H.; MODLIN, IRVIN M. Archives of Surgery April, 1988; 123: 431-4331988;

LINE COUNT: 00158 WORD COUNT: 02190

...might yield better results than prostaglandin therapy alone. It may also be useful in future clinical trials of the prostaglandins to monitor pepsinogen secretion via serum pepsinogen levels or mucosal biopsies.

Changes in pepsinogen secretion should be considered in the choice...

6/3,K/38 (Item 10 from file: 442)

DIALOG(R) File 442:AMA Journals

(c) 2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00039740

Copyright (C) 1986 American Medical Association

Cefoxitin Plus Tobramycin and Clindamycin Plus Tobramycin; A Prospective Randomized Comparison in the Therapy of Mixed Aerobic/Anaerobic Infections (ORIGINAL ARTICLE)

NICOLLE, LINDSAY E.; HARDING, GODFREY K. M.; LOUIE, THOMAS J.; THOMSON, MARGARET J.; BLANCHARD, ROBERT J.

Archives of Surgery

August, 1986; 121: 891-8961986;

LINE COUNT: 00359 WORD COUNT: 04960

...surgeon and the infectious disease consultant.

Prior to institution of therapy with the study antimicrobial, blood cultures, hematologic and biochemistry studies, and urine specimens were obtained. Patients receiving the study antimicrobial were monitored for clinical response, number of daily bowel movements biochemistry, hematology, urinalysis, and serum tobramycin levles, and repeated cultures when appropriate. Tests were repeated at the termination of the...

6/3,K/39 (Item 11 from file: 442)

DIALOG(R) File 442:AMA Journals (c) 2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00031656

Copyright (C) 1986 American Medical Association

Total and Free Serum Concentrations of Carbamazepine and Carbamazepine-10,11-Epoxide in Children With Epilepsy (ORIGINAL CONTRIBUTION)

AGBATO, OLATOYE A.; ELYAS, ABDULMAJID A.; PATSALOS, PHILIP N.; BRETT, EDWARD M.; LASCELLES, PETER T.

Archives of Neurology

November, 1986; 43: 1111-11161986;

LINE COUNT: 00326 WORD COUNT: 04502

CITED REFERENCES:

- ...6. Hoppener RJ, Kuyer A, Meyer JWA, et al: Correlation between daily fluctuations of carbamazepine **serum** levels and intermittent side effects. Epilepsia 1980;21:341-350.
- 7. Perucca E: Free level monitoring of antiepileptic drugs: Clinical usefulness and case studies. Clin Pharmacokinet 1984;9(suppl 1):7178.
- 8. Bourgeois BFD, Wad N: Individual and combined...

6/3,K/40 (Item 12 from file: 442)

DIALOG(R) File 442:AMA Journals

(c) 2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00029991 Copyright (C) 1987 American Medical Association

Ambulatory Blood Pressure Monitoring in Patients With Panic Disorder (ORIGINAL INVESTIGATIONS)

WHITE, WILLIAM B.

Archives of Internal Medicine

November, 1987; 147: 1973-19751987;

LINE COUNT: 00159 WORD COUNT: 02199

... studies, (Ref. 4) and increases of ten to 35 bpm have been noted using ambulatory monitoring equipment in clinical studies. (Ref. 5,6) Recently, we described two patients who presented with episodic elevations of the blood pressure secondary to panic disorder. (Ref. 7) These cases made us recognize that patients of... not changed or have increased minimally. (Ref. 20,21)

It is apparent from this first clinical study of blood pressure monitoring in patients with panic disorder that further cardiovascular studies are desirable. Evaluation for the presence...

Studios are appropriately

6/3,K/41 (Item 13 from file: 442)

DIALOG(R) File 442: AMA Journals

(c)2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00027056

Copyright (C) 1984 American Medical Association

Tricyclic Antidepressant Therapy for Peptic Ulcer Disease (TODAY'S PRACTICE OF MEDICINE)

RIES, RICHARD K.; GILBERT, DAVID A.; KATON, WAYNE

Archives of Internal Medicine March, 1984; 144: 566-5691984;

LINE COUNT: 00290 WORD COUNT: 04010

... Other advantages of these agents include their long half-lives, low cost, and readily available **serum monitoring**. Further **clinical studies** with detailed physiologic and psychologic observations and **serum** monitoring using TCAs in patients with peptic ulcer disease are needed. (Arch Intern Med 1984...

6/3,K/42 (Item 14 from file: 442)

DIALOG(R) File 442:AMA Journals

(c) 2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00026762

Copyright (C) 1987 American Medical Association

Imipramine in Prepubertal Major Depressive Disorders (ORIGINAL ARTICLE)

PUIG-ANTICH, JOAQUIM; PEREL, JAMES M.; LUPATKIN, WILLIAM; CHAMBERS, WILLIAM J.; TABRIZI, MARY ANN; KING, JANET; GOETZ, RAYMOND; DAVIES, MARK; STILLER, RICHARD L.

Archives of General Psychiatry

January, 1987; 44: 81-891987;

LINE COUNT: 00606 WORD COUNT: 08376

... of the pills (except the pharmacist) and to plasma levels (except the pharmacology laboratory). The **clinical monitor** and the **research** nurse collected side effect scales, ECGs, and **blood** pressure values. At the end of the imipramine protocol, the **clinical monitor** and the **research** nurse jointly guessed the patient's assignment to drug or placebo. Although they were not...

...QRS width. This eventuality had been anticipated before the start of the protocol. Thus, ECGs, blood pressure values, and side effect scales were kept from all clinical personnel except the clinical monitor and research nurse, who obtained such measurements and used them in monitoring dosage and systematically kept them...

6/3,K/43 (Item 15 from file: 442)

DIALOG(R) File 442: AMA Journals

(c) 2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00023312

Copyright (C) 1989 American Medical Association

AIDS Antigen Test (FROM THE FOOD AND DRUG ADMINISTRATION)

NIGHTINGALE, BY STUART L.

JAMA, The Journal of the American Medical Association

December 15, 1989; 262: 32541989;

LINE COUNT: 00038 WORD COUNT: 00527

... have been reported are monitoring the presence of HIV-1 antigen in cocultures of peripheral **blood** lymphocytes and, in **clinical research**, **monitoring** treatment of HIV-1-infected individuals with antiviral drugs. The test is not intended as...

6/3,K/44 (Item 16 from file: 442)

DIALOG(R) File 442: AMA Journals

(c) 2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00020656

Copyright (C) 1989 American Medical Association

Average Daily Blood Pressure, Not Office Blood Pressure, Determines Cardiac Function in Patients With Hypertension (ORIGINAL CONTRIBUTIONS)

WHITE, WILLIAM B.; SCHULMAN, PETER; MCCABE, ELLEN J.; DEY, HOLLEY M.

JAMA, The Journal of the American Medical Association

February 10, 1989; 261: 873-8771989;

LINE COUNT: 00265 WORD COUNT: 03667

...in the Hypertension Unit at the University of Connecticut Health Center, Farmington, who underwent ambulatory blood pressure monitoring for clinical or pharmacologic research purposes. Over a period of four years (1983 through 1987), 720 patients underwent ambulatory blood pressure monitoring and 77 of these patients were never previously treated with antihypertensive therapy. Twenty...

6/3,K/45 (Item 17 from file: 442)

DIALOG(R) File 442:AMA Journals

(c) 2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00020025 Copyright (C) 1988 American Medical Association

Adverse Events Associated With 24-Hour Ambulatory Sphygmomanometry (LETTERS)

BURRIS, JAMES F.

JAMA, The Journal of the American Medical Association

November 4, 1988; 260: 2508-25091988;

LINE COUNT: 00051 WORD COUNT: 00705

...an ambulatory blood pressure monitoring device (Table).

Adverse Events Associated With Use of an Ambulatory **Blood** Pressure Monitoring Device

(SEE ORIGINAL SOURCE)

Each of these patients underwent ambulatory **blood** pressure **monitoring** as part of a **clinical trial** of an investigational antihypertensive drug. The devices had been tested, applied, and activated by clinical...

6/3,K/46 (Item 18 from file: 442)

DIALOG(R) File 442: AMA Journals (c) 2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00016665

Copyright (C) 1987 American Medical Association

Phenylpropanolamine in Patients With Autonomic Impairment (LETTERS)

KROENKE, MAJ KURT

JAMA, The Journal of the American Medical Association

December 4, 1987; 258: 3121-31221987;

LINE COUNT: 00101 WORD COUNT: 01405

... the "Methods" section, one can picture the 14 study patients with autonomic dysfunction, hooked to monitors in a clinical research center, nervously swallowing pills they are told might raise their blood pressure. Indeed, the principles of informed consent would compel the investigators to disclose even more...

6/3,K/47 (Item 19 from file: 442)

DIALOG(R)File 442:AMA Journals

(c) 2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00014377

Copyright (C) 1987 American Medical Association

Nutritional Therapy for High Blood Pressure; Final Report of a Four-Year Randomized Controlled Trial -- The Hypertension Control Program (ORIGINAL CONTRIBUTIONS)

STAMLER, ROSE; STAMLER, JEREMIAH; GRIMM, RICHARD; GOSCH, FLORA C.; ELMER, PATRICIA; DYER, ALAN; BERMAN, REUBEN; FISHMAN, JOAN; VAN HEEL, NANCY; CIVINELLI, JEAN; MCDONALD, ARLINE JAMA, The Journal of the American Medical Association March 20, 1987; 257: 1484-14911987;

LINE COUNT: 00488 WORD COUNT: 06742

... blood pressure and levels of biochemical variables affected by the drugs. Length of observation and sample size did not permit utilizing morbidity and mortality as trial end points, although these were monitored.

Sample Size

Sample size was calculated to demonstrate (with alpha 0.05 and beta 0.10) the following...

(Item 20 from file: 442) 6/3,K/48

DIALOG(R) File 442:AMA Journals

(c)2003 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00005379

Copyright (C) 1984 American Medical Association

The Lipid Research Clinics Coronary Primary Prevention Trial Results; I. Reduction in Incidence of Coronary Heart Disease (ORIGINAL CONTRIBUTIONS)

PROGRAM, LIPID RESEARCH CLINICS

JAMA, The Journal of the American Medical Association

January 20, 1984; 251: 351-3641984;

LINE COUNT: 00809 WORD COUNT: 11169

... Little or no differences were seen between the two treatment groups for most of the clinical chemical tests monitored during the study (Appendix C). During the first year, serum alkaline phosphatase levels, iron binding capacity, SGOT levels, and the WBC count were higher in...

(Item 1 from file: 149) 6/3,K/49

DIALOG(R)File 149:TGG Health&Wellness DB(SM)

(c) 2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 63815799 Linking Community-Based Blood Pressure Measurement to Clinical Care: A Randomized Controlled Trial of Outreach and Tracking by Community

Health Workers.

Krieger, James; Collier, Cheza; Song, Lin; Martin, Donald

The American Journal of Public Health, 89, 6, 856

June,

1999

ISSN: 0090-0036 PUBLICATION FORMAT: Magazine/Journal; Refereed

LANGUAGE: English RECORD TYPE: Abstract TARGET AUDIENCE: Academic

Linking Community-Based Blood Pressure Measurement to Clinical Care: A Randomized Controlled Trial of Outreach and Tracking by Community Health Workers.

(Item 2 from file: 149) 6/3,K/50

DIALOG(R)File 149:TGG Health&Wellness DB(SM)

(c) 2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 57943233

There's no place like home: distance technologies of patient

monitoring. (The Impact of New Technologies in Medicine)

British Medical Journal, 319, 7220, 1309 Nov 13,

1999

PUBLICATION FORMAT: Magazine/Journal ISSN: 0959-8146 LANGUAGE: English

RECORD TYPE: Fulltext TARGET AUDIENCE: Professional

WORD COUNT: 269 · LINE COUNT: 00028

meaningful communication and decision support systems are also needed for patients and clinicians. Several randomised clinical trials of computerised home glucose monitoring found improved outcomes in diabetes care. Trials of home monitoring of blood pressure show enhanced efficiency. Failure to improve outcomes through home monitoring of high-risk pregnancies...

6/3,K/51 (Item 3 from file: 149)

DIALOG(R) File 149:TGG Health & Wellness DB(SM)

(c) 2003 The Gale Group. All rts. reserv.

01856471 SUPPLIER NUMBER: 55807393 (USE FORMAT 7 OR 9 FOR FULL TEXT)
Colour doppler imaging for diagnosis of intracranial
hypotension. (Statistical Data Included)

Chao-Ching, Chen; Cheau-Lin, Luo; Shuu-Jiun, Wang; Chang-Ming, Chern;

Jong-Ling, Fuh; Sheng-Han, Lin; Han-Hwa, Hu

The Lancet, 354, 9181, 826

Sept 4,

1999

DOCUMENT TYPE: Statistical Data Included PUBLICATION FORMAT: Magazine/Journal; Refereed ISSN: 0099-5355 LANGUAGE: English

RECORD TYPE: Fulltext TARGET AUDIENCE: Professional

WORD COUNT: 3323 LINE COUNT: 00314

... Moreover, since changes in the blood flow in the superior ophthalmic vein were reversible after **blood** -patch treatment, CDFI would also be useful in **monitoring** the therapeutic effect of **clinical** management. However, this preliminary **study** included only a small number of participants, so further investigations in a larger population are...

6/3,K/52 (Item 4 from file: 149)

DIALOG(R) File 149:TGG Health&Wellness DB(SM)

(c) 2003 The Gale Group. All rts. reserv.

01813155 SUPPLIER NUMBER: 53520315 (USE FORMAT 7 OR 9 FOR FULL TEXT)

What's new in hypertension: update on angiotensin II receptor

blockers. (includes related article on the role of angiotensin II in the development of hypertension)

Black, Henry R.

Consultant, 38, 11, 2715(7)

Nov,

1998

PUBLICATION FORMAT: Magazine/Journal; Refereed ISSN: 0010-7069 LANGUAGE: English RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE:

Professional

WORD COUNT: 3491 LINE COUNT: 00308

... receptor in vitro than losartan is, but it is unclear whether this feature has any clinical implications. (15)

EFFICACY

Clinical studies using ambulatory blood pressure monitoring

have shown that all the ARBs produce smooth antihypertensive actions over a 24-hour period...

6/3,K/53 (Item 5 from file: 149)
DIALOG(R)File 149:TGG Health&Wellness DB(SM)
(c) 2003 The Gale Group. All rts. reserv.

01739991 SUPPLIER NUMBER: 20062550 (USE FORMAT 7 OR 9 FOR FULL TEXT)
Update on rifampin drug interactions. (part 3)

Strayhorn, Valerie A.; Baciewicz, Anne M.; Self, Timothy H. Archives of Internal Medicine, v157, n21, p2453(6)

Archives of Internal Medicine, v15/, n21, p2453(6) Nov 24,

Nov 24, 1997

PUBLICATION FORMAT: Magazine/Journal; Refereed ISSN: 0003-9926 LANGUAGE: English RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE:

Professional WORD COUNT: 4671 LINE COUNT: 00489

... clinical response;

blocking agents increased propanolol or metoprolol

dose may be needed

Chloramphenicol((dagger)) Monitor serum chloramphenicol

concentrations: may need to increase

dosage

Dapsone Monitor clinical response: additional

study needed when used for
Pneumocystis carinii prophylaxis;
monitor for hematologic toxic effects

Diazepam((dagger)) Monitor...

6/3,K/54 (Item 6 from file: 149)
DIALOG(R)File 149:TGG Health&Wellness DB(SM)
(c) 2003 The Gale Group. All rts. reserv.

01730722 SUPPLIER NUMBER: 19987821 (USE FORMAT 7 OR 9 FOR FULL TEXT)
Stress management and exercise training in cardiac patients with myocardial ischemia: effects on prognosis and evaluation of mechanisms.

Blumenthal, James A.; Jiang, Wei; Babyak, Michael A.; Krantz, David S.; Frid, David J.; Coleman, R. Edward; Waugh, Robert; Hanson, Michael; Appelbaum, Mark; O'Connor, Christopher; Morris, James J. Archives of Internal Medicine, v157, n19, p2213(11) Oct 27,

1997

PUBLICATION FORMAT: Magazine/Journal; Refereed ISSN: 0003-9926
LANGUAGE: English RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE:
Professional

WORD COUNT: 8022 LINE COUNT: 00699

not participation. Of the 136 patients initially enrolled in the study, 107 met the criteria of **exhibiting** myocardial ischemia with either mental stress testing or ambulatory **monitoring** and were included in the **clinical trial**. All subjects received their group assignments immediately following the completion of the pretreatment evaluation. Those ...

6/3,K/55 (Item 7 from file: 149)
DIALOG(R)File 149:TGG Health&Wellness DB(SM)

(c) 2003 The Gale Group. All rts. reserv.

01720306 SUPPLIER NUMBER: 19802771 (USE FORMAT 7 OR 9 FOR FULL TEXT)
The history of multiple sclerosis. (part 2)

Rolak, Loren A.

Inside MS, v14, n3, p14(2)

Fall,

1996

PUBLICATION FORMAT: Newsletter ISSN: 0739-9774 LANGUAGE: English

RECORD TYPE: Fulltext TARGET AUDIENCE: Consumer

WORD COUNT: 986 LINE COUNT: 00087

... treated people with MS in the 1950s continued to suspect the cause lay in impaired blood flow -- so circulation stimulators dominated treatment. These therapies were used without controlled studies to track the results, called clinical trials, so no reproducible or valid information could emerge about their effectiveness.

Breakthroughs expand knowledge but...

6/3,K/56 (Item 8 from file: 149)

DIALOG(R) File 149: TGG Health & Wellness DB(SM)

(c) 2003 The Gale Group. All rts. reserv.

01717679 SUPPLIER NUMBER: 19798382 (USE FORMAT 7 OR 9 FOR FULL TEXT) Randomised trial of single and repeated fibrin glue compared with injection of polidocanol in treatment of bleeding peptic ulcer.

Rutgeerts, Paul; Rauws, Erik; Wara, Pal; Swain, Paul; Hoos, Anton; Solleder, Eva; Halttunen, Jorma; Dobrilla, Giorgio; Richter, Gerhard; Prassler, Roland

The Lancet, v350, n9079, p692(5)

Sep 6,

1997

PUBLICATION FORMAT: Magazine/Journal; Refereed ISSN: 0099-5355
LANGUAGE: English RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE:

Professional

WORD COUNT: 4090 LINE COUNT: 00356

... with the Declaration of Helsinki. Data were recorded carefully in case-record forms and the **study** was **monitored** according to good **clinical** practice regulations.33

Statistics The **sample** size of 250 people per treatment group was based on an expected recurrent bleeding rate...

6/3,K/57 (Item 9 from file: 149)

DIALOG(R) File 149: TGG Health & Wellness DB(SM)

(c) 2003 The Gale Group. All rts. reserv.

01689291 SUPPLIER NUMBER: 19367523 (USE FORMAT 7 OR 9 FOR FULL TEXT)

New drug for type II: troglitazone increases insulin sensitivity.

(Rezulin) (includes related information)

McCarren, Marie

Diabetes Forecast, v50, n5, p26(4)

May,

1997

PUBLICATION FORMAT: Magazine/Journal ISSN: 0095-8301 LANGUAGE: English

RECORD TYPE: Fulltext TARGET AUDIENCE: Consumer

WORD COUNT: 1777 LINE COUNT: 00139

... at the University of Chicago, has had about 20 patients on troglitazone as part of **clinical studies**. He recommends that people **monitor** their **blood** glucose levels three or four times a day and keep in close contact with their...

6/3,K/58 (Item 10 from file: 149)
DIALOG(R)File 149:TGG Health&Wellness DB(SM)
(c) 2003 The Gale Group. All rts. reserv.

01675093 SUPPLIER NUMBER: 19156192 (USE FORMAT 7 OR 9 FOR FULL TEXT) Hyperventilation and head injury: controversies and concerns.

Geraci, Eunell B.; Geraci, Thomas A.

Journal of Neuroscience Nursing, v28, n6, p381(7)

Dec,

1996

PUBLICATION FORMAT: Magazine/Journal ISSN: 0888-0395 LANGUAGE: English

RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE: Professional

WORD COUNT: 5183 LINE COUNT: 00453

...ABSTRACT: standard treatment for serious head injuries for over 20 years. Recent technological advances in patient **monitoring** and related **medical research** suggest that unmonitored hyperventilation can lead to secondary brain injury, such as ischemia, by reducing **blood** flow during the prehospital and early emergency phases of treatment, making its continued usage controversial...

6/3,K/59 (Item 11 from file: 149)
DIALOG(R)File 149:TGG Health&Wellness DB(SM)
(c) 2003 The Gale Group. All rts. reserv.

01669810 SUPPLIER NUMBER: 19016982 (USE FORMAT 7 OR 9 FOR FULL TEXT)
Oat bran concentrate bread products improve long-term control of diabetes:
a pilot study.

Pick, Mary E.; Hawrysh, Zenia J.; Gee, Margaret I.; Toth, Ellen; Garg,

Manohar L.; Hardin, Robert T.

Journal of the American Dietetic Association, v96, n12, p1254(8)

Dec,

1996

PUBLICATION FORMAT: Magazine/Journal ISSN: 0002-8223 LANGUAGE: English

RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE: Professional

WORD COUNT: 6187 LINE COUNT: 00534

... products as bread exchanges. Daily energy intake was individualized for weight maintenance.

Dietary intakes were **monitored** by a **research** dietitian; a physician provided **medical** management. Patients were seen every 3 weeks to make anthropometric and dietary assessments, to discuss...

6/3,K/60 (Item 12 from file: 149)
DIALOG(R)File 149:TGG Health&Wellness DB(SM)
(c) 2003 The Gale Group. All rts. reserv.

O1653801 SUPPLIER NUMBER: 17847456 (USE FORMAT 7 OR 9 FOR FULL TEXT)
Nutrition intervention program of the Modification of Diet in Renal Disease
Study: a self-management approach. (includes related article)
Gillis, Bonnie P.; Caggiula, Arlene W.; Chiavacci, Anne T.; Coyne, Terry;
Doroshenko, Linda; Milas, N. Carole; Nowalk, Mary Patricia; Scherch, Laura

Journal of the American Dietetic Association, v95, n11, p1288(7) 1995

ISSN: 0002-8223 LANGUAGE: English PUBLICATION FORMAT: Magazine/Journal

RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE: Professional

LINE COUNT: 00489 WORD COUNT: 5679

and guidelines for vegetarian meal planning. A few participants were given several days of actual sample meals or were admitted to a research center where food intake was monitored closely; these activities provided insights for the interpretation and improvement of estimations of protein intake...

(Item 13 from file: 149) 6/3,K/61 DIALOG(R) File 149:TGG Health & Wellness DB(SM) (c) 2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 18319731 01619849 Forecasting Alzheimer's disease: brain scans and writing samples may predict dementia. (Cover Story)

Fackelmann, Kathleen Science News, v149, n20, p312(2) May 18,

1996

PUBLICATION FORMAT: Magazine/Journal DOCUMENT TYPE: Cover Story

0036-8423 LANGUAGE: English RECORD TYPE: Fulltext; Abstract

TARGET AUDIENCE: Academic; Consumer

LINE COUNT: 00165 WORD COUNT: 2136

s disease," he says. First, however, scientists must determine whether the combination of PET and blood tests really does forecast dementia, comments Mony J. de Leon, an Alzheim-er's researcher at New York University Medical Center. To do this, researchers must continue monitoring their volunteers to see whether they actually develop Alzheimer's, he adds.

This widespread scientific...

(Item 14 from file: 149) 6/3,K/62 DIALOG(R) File 149:TGG Health & Wellness DB(SM) (c) 2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 17737854 01607279 Reach new sales heights: dealers share their merchandising tips for blood pressure monitors.

Nones, Rachelle Independent Living Provider, v10, n7, p60(2) Nov-Dec,

1995 PUBLICATION FORMAT: Magazine/Journal LANGUAGE: English RECORD TYPE:

Fulltext TARGET AUDIENCE: Consumer LINE COUNT: 00080 WORD COUNT: 982

In the mid-1980s, Kaiser Permanente Medical Care Program in San Francisco conducted a home blood pressure monitoring clinical study of 467 patients. Results showed that subjects who took their pressures at home made 1...

(Item 15 from file: 149) 6/3,K/63 DIALOG(R)File 149:TGG Health&Wellness DB(SM) (c) 2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 17467906 Antithrombotic therapy in children. (includes related article) (Fourth Annual American College of Chest Physicians Consensus Conference on Antithrombotic Therapy, April 1995, Tucson, Arizona)

Michelson, Alan D.; Bovill, Edward; Andrew, Maureen

Chest, v108, n4, p506S(17)

Oct,

1995 ISSN: 0012-3692 LANGUAGE: English PUBLICATION FORMAT: Magazine/Journal

RECORD TYPE: Fulltext TARGET AUDIENCE: Professional

LINE COUNT: 01211 13382 WORD COUNT:

solutions for optimizing therapy with oral anticoagulants in children include pediatric anticoagulation clinics, [1] whole blood PT/INR monitors used at home, [2] and clinical trials to determine if lower, safer INR ranges are as efficacious.

Adverse Effects of Oral Anticoagulants...

(Item 16 from file: 149) 6/3,K/64

DIALOG(R) File 149:TGG Health&Wellness DB(SM) (c) 2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 15949341 trials with ambulatory blood pressure monitoring : fewer 01496911 Clinical patients needed?

Staessen, Jan A.; Thijs, Lutgarde; Mancia, Giuseppe; Parati, Gianfranco; O'Brien, Eoin T.

The Lancet, v344, n8936, p1552(5)

Dec 3,

1994 ISSN: 0099-5355 LANGUAGE: English PUBLICATION FORMAT: Magazine/Journal

RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE: Professional

LINE COUNT: 00253 3093 WORD COUNT:

trials with ambulatory blood pressure monitoring : fewer 、 Clinical patients needed?

pressure reduction at a given time (eg, the time of peak plasma concentration) and the **blood** pressure changes at all other times.

It has also been suggested that ambulatory monitoring may trials by the early identification facilitate the conduct of clinical and exclusion of subjects whose blood pressure is raised only in the clinic environment (white coat reactors).[19,26] This feature...

(Item 17 from file: 149) 6/3,K/65

DIALOG(R) File 149:TGG Health & Wellness DB(SM)

(c) 2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 15928000 Long-term xanthine therapy of asthma: enprofylline and theophylline 01495895 compared.

Chapman, Kenneth R.; Ljungholm, Karin; Kallen, Anders Chest, v106, n5, p1407(7) Nov,

1994

PUBLICATION FORMAT: Magazine/Journal ISSN: 0012-3692 LANGUAGE: English

RECORD TYPE: Fulltext TARGET AUDIENCE: Professional

WORD COUNT: 4765 LINE COUNT: 00412

... level stability in long-term therapy would call into question the need for routine, frequent **blood** level **monitoring**. Such a hypothesis deserves further **clinical study**.

Recent publications have suggested that long-term bronchodilator administration to patients with asthma may lead...

6/3,K/66 (Item 18 from file: 149)

DIALOG(R)File 149:TGG Health&Wellness DB(SM)

(c) 2003 The Gale Group. All rts. reserv.

01487733 SUPPLIER NUMBER: 15739728 (USE FORMAT 7 OR 9 FOR FULL TEXT)
Pharmacological treatment of hypertension. (Hypertension Octet)

Swales, J.D.

The Lancet, v344, n8919, p380(6)

August 6,

1994

PUBLICATION FORMAT: Magazine/Journal ISSN: 0099-5355 LANGUAGE: English

RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE: Professional

WORD COUNT: 5619 LINE COUNT: 00469

function in patients with hypertension. JAMA 1989; 261: 873-77. [24] Mancia G. Ambulatory blood **pressure** monitoring: **research** and clinical applications. J Hypertens 1990; 8 (suppl 7): s1-13. [25] Alderman MH. Blood **pressure** management: individualised treatment based on absolute risk and the potential for benefit. Ann Intern Med...

6/3,K/67 (Item 19 from file: 149)

DIALOG(R) File 149:TGG Health&Wellness DB(SM)

(c) 2003 The Gale Group. All rts. reserv.

01483989 SUPPLIER NUMBER: 15596744 (USE FORMAT 7 OR 9 FOR FULL TEXT)

Status of Immupath reported as potential AIDS treatment.

AIDS Weekly, p8(2)

July 18,

1994

DOCUMENT TYPE: evaluation PUBLICATION FORMAT: Newsletter ISSN:

1069-1456 LANGUAGE: English RECORD TYPE: Fulltext TARGET AUDIENCE:

Professional

WORD COUNT: 425 LINE COUNT: 00042

... 3). (Decreasing CD4 cell concentrations are indicators of AIDS

progression.)

The FDA has approved the **study** 's design, **sample** size, endpoints, **clinical monitoring** and manufacturing plan but has questioned the viral inactivation procedures used in processing Immupath. On...

6/3,K/68 (Item 20 from file: 149)

DIALOG(R)File 149:TGG Health&Wellness DB(SM)

(c) 2003 The Gale Group. All rts. reserv.

01483303 SUPPLIER NUMBER: 15505098 (USE FORMAT 7 OR 9 FOR FULL TEXT) Problems in clinical trials go far beyond misconduct. (includes glossary

and related articles on effects of ignorance and on meta-analysis and megatrials) (Special News Report: A Close Look at Clinical Trials)

Nowak, Rachel

Science, v264, n5165, p1538(4)

June 10,

1994

ISSN: 0036-8075 LANGUAGE: English PUBLICATION FORMAT: Magazine/Journal

RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE: Academic

LINE COUNT: 00355 4314 WORD COUNT:

for death. As a result, many clinical trials look for effects on CD4 counts and serum cholesterol levels rather than monitoring hard clinical endpoints. Most experts in clinical trials agree that it's fine to use surrogate markers to identify promising new agents in...

(Item 21 from file: 149) 6/3, K/69

DIALOG(R) File 149:TGG Health & Wellness DB(SM)

(c) 2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 15518780 01482795 HemaCare reports on regulatory status of Immupath.

AIDS Weekly, p14(1)

June 27,

PUBLICATION FORMAT: Newsletter ISSN: 1069-1456 LANGUAGE: English

RECORD TYPE: Fulltext TARGET AUDIENCE: Professional

LINE COUNT: 00042 WORD COUNT: 429

lymphocyte concentrations of 50-200/cells per [mm.sup.3]. The FDA has approved the study 's design, sample size, endpoints, monitoring and manufacturing plan but has questioned the viral inactivation procedures used in processing Immupath. On...

(Item 22 from file: 149) 6/3,K/70

DIALOG(R) File 149:TGG Health&Wellness DB(SM)

(c) 2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 15231319 Hepatotoxic effects of tacrine administration in patients with Alzheimer's 01479406

Watkins, Paul B.; Zimmerman, Hyman J.; Knapp, Margaret J.; Gracon, Stephen disease.

I.; Lewis, Karen W. JAMA, The Journal of the American Medical Association, v271, n13, p992(7) April 6,

1994 ISSN: 0098-7484 LANGUAGE: English PUBLICATION FORMAT: Magazine/Journal

RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE: Professional

LINE COUNT: 00522 6346 WORD COUNT:

TEXT:

...seven of the 50 patients enrolled.[5] As a result of these initial studies have required frequent observations, subsequent clinical (weekly) monitoring of serum ALT levels.

(Item 23 from file: 149) 6/3,K/71

DIALOG(R) File 149:TGG Health&Wellness DB(SM)

(c) 2003 The Gale Group. All rts. reserv.

01435184 SUPPLIER NUMBER: 15024488 (USE FORMAT 7 OR 9 FOR FULL TEXT)
1993 guidelines for the management of mild hypertension: memorandum from a
WHO/ISH meeting. (World Health Organization, International Society of
Hypertension)

Bulletin of the World Health Organization, v71, n5, p503(15) Sept-Oct,

1993

PUBLICATION FORMAT: Magazine/Journal ISSN: 0042-9686 LANGUAGE: English RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE: Professional WORD COUNT: 10532 LINE COUNT: 00908

... a follow-up. Clin. exper. hypertens., A14: 45-53 (1992). [12] Mancia, G. Ambulatory blood **pressure** monitoring: **research** and clinical **applications**. J. hypertens., 8 (suppl. 7): S1-S13 (1990). [13] O'Brien, E. et al. The...

6/3,K/72 (Item 24 from file: 149)
DIALOG(R)File 149:TGG Health&Wellness DB(SM)
(c) 2003 The Gale Group. All rts. reserv.

01434803 SUPPLIER NUMBER: 14742164 (USE FORMAT 7 OR 9 FOR FULL TEXT)
Summary of 1993 World Health Organisation-International Society of
Hypertension guidelines for the management of mild hypertension.
British Medical Journal, v307, n6918, p1541(6)

Dec 11,

1993
PUBLICATION FORMAT: Magazine/Journal ISSN: 0959-8146 LANGUAGE: English
RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE: Professional
WORD COUNT: 4509 LINE COUNT: 00453

... a follow-up. Clin Exp Hypertens [A] 1992;A14:45-53.

[12] Mancia G. Ambulatory blood pressure monitoring: research and clinical applications. J Hypertens 1990;8(suppl 7):S1-S13.

[13] Zanchetti A. What **blood** pressure level should be treated? In: Laragh JH, Brenner BM, eds. Hypertension: pathophysiology, diagnosis, and

6/3,K/73 (Item 25 from file: 149)
DIALOG(R)File 149:TGG Health&Wellness DB(SM)
(c) 2003 The Gale Group. All rts. reserv.

01434178 SUPPLIER NUMBER: 14386838 (USE FORMAT 7 OR 9 FOR FULL TEXT)

Dexamethasone therapy for bacterial meningitis in children. (Swiss

Meningitis Study Group)

Schaad, Urs B.; Lips, Ueli; Gnehm, Hanspeter E.; Blumberg, Avihay; Heinzer, Ivo; Wedgwood, Joanna

The Lancet, v342, n8869, p457(5)

August 21,

1993

PUBLICATION FORMAT: Magazine/Journal ISSN: 0099-5355 LANGUAGE: English RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE: Professional WORD COUNT: 4116 LINE COUNT: 00345

... and at the end of antibiotic therapy.[17,18] Adverse or toxic reactions to the **study** drugs were sought by daily **clinical** observation and laboratory **monitoring** of complete **blood** cell count, electrolytes, creatinine, and liver enzymes in **serum**, glucose in urine, and occult

blood in stool, on days 1,2, and 3, and...

(Item 26 from file: 149) 6/3,K/74 DIALOG(R) File 149:TGG Health&Wellness DB(SM) (c) 2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 13269131 01425803 IND granted for immune reconstitution therapy. (Investigational New Drug application by Health Professionals Inc.'s Center for Special Immunology) (U.S. Food and Drug Administration)

AIDS Weekly, v36, n34, p6(1)

Sept 6,

1993 PUBLICATION FORMAT: Newsletter ISSN: 1069-1456 LANGUAGE: English

RECORD TYPE: Fulltext TARGET AUDIENCE: Professional

LINE COUNT: 00018 199 WORD COUNT:

it with the FDA within six weeks. The expanded program will be administered as a monitored clinical trial," Reiter concluded. CSI's immune reconstitution therapy includes an infusion of peripheral blood lymphocytes, which are specific types of white blood cells deficient in patients with AIDS. An...

(Item 27 from file: 149) 6/3,K/75 DIALOG(R) File 149:TGG Health&Wellness DB(SM) (c) 2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 12280406 Longitudinal evaluation of prostate-specific antigen levels in men with and without prostate disease.

Carter, H. Ballentine; Pearson, Jay D.; Metter, E. Jeffrey; Brant, Larry J. ; Chan, Daniel W.; Andres, Reubin; Fozard, James L.; Walsh, Patrick C. JAMA, The Journal of the American Medical Association, v267, n16, p2215(6) April 22,

1992

ISSN: 0098-7484 LANGUAGE: English PUBLICATION FORMAT: Magazine/Journal RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE: Professional LINE COUNT: 00365 4402 WORD COUNT:

case-control study may not be representative of changes that occur in all men being monitored by PSA levels in clinical practice. In this study , PSA measurements were performed on stored serum samples and not on consecutively collected samples over many years as would be the case in clinical practice where the interassay variation...

(Item 28 from file: 149) 6/3,K/76 DIALOG(R) File 149:TGG Health&Wellness DB(SM) (c) 2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 11951693 01351688 Measurement of isometric force in children with and without Duchenne's muscular dystrophy.

Brussock, Christine M.; Haley, Stephen M.; Munsat, Theodore L.; Bernhardt,

Physical Therapy, v72, n2, p105(10)

Feb, 1992

Bode Akintola 25-Apr-03

PUBLICATION FORMAT: Magazine/Journal ISSN: 0031-9023 LANGUAGE: English RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE: Professional WORD COUNT: 4225 LINE COUNT: 00475

markers that accurately describe disease progression in individuals with DMD have been reported. Levels of **serum** creatine phosphokinase and 3-methylhistidine have been **monitored** in **clinical trials**, but they have been ineffective in judging therapeutic responses to new interventions. [3,4]

The...

6/3,K/77 (Item 29 from file: 149)
DIALOG(R)File 149:TGG Health&Wellness DB(SM)
(c) 2003 The Gale Group. All rts. reserv.

O1314060 SUPPLIER NUMBER: 11986669 (USE FORMAT 7 OR 9 FOR FULL TEXT)
'White coat' hypertension in children.
Hornsby, J. Larry; Mongan, Patrick F.; Taylor, A. Thomas; Treiber, Frank A.
Journal of Family Practice, v33, n6, p617(7)
Dec,
1991

PUBLICATION FORMAT: Magazine/Journal ISSN: 0094-3509 LANGUAGE: English RECORD TYPE: Fulltext TARGET AUDIENCE: Professional WORD COUNT: 3463 LINE COUNT: 00342

- 24-hour blood pressure readings in the normal range) also limits accurate diagnosis when evaluating **blood** pressure in the **medical** setting. [3] **Research** with ambulatory **blood** pressure **monitoring** (ABPM) clearly demonstrates the importance of white coat hypertension in making an accurate diagnosis of...dimensions of "office hypertension." Behav Res Ther 1985; 23:609-12.
- [8] Mancia G. Ambulatory **blood** pressure **monitoring** in hypertension **research** and **clinical** practice. ISH Hypertension Yearbook 1986; 93-115.
 - [9] Waeber B, Scherrer U, Petrilo A, et...

6/3,K/78 (Item 30 from file: 149)
DIALOG(R)File 149:TGG Health&Wellness DB(SM)
(c) 2003 The Gale Group. All rts. reserv.

01301693 SUPPLIER NUMBER: 10975717 (USE FORMAT 7 OR 9 FOR FULL TEXT)

Ambulatory blood pressure monitoring in hypertension.

Hornsby, J. Larry; Mongan, Patrick F.; Taylor, A. Thomas

American Family Physician, v43, n5, p1631(8)

May,
1991

PUBLICATION FORMAT: Magazine/Journal ISSN: 0002-838X LANGUAGE: English RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE: Professional WORD COUNT: 2750 LINE COUNT: 00291

... to the development of ABPM monitors that provide an accurate and reliable method for assessing **blood** pressure for **clinical** as well as **research** purposes. (26-28)

MEASUREMENT INTERVALS

Monitors may be programmed to sample blood pressure at various intervals, usually every five to 60 minutes for 24 to 48 hours...

6/3,K/79 (Item 31 from file: 149)
DIALOG(R)File 149:TGG Health&Wellness DB(SM)
(c) 2003 The Gale Group. All rts. reserv.

01293059 SUPPLIER NUMBER: 10346599 (USE FORMAT 7 OR 9 FOR FULL TEXT)
Management of mild hypertension: selecting an antihypertensive regime.

Perez-Stable, Eliseo J.

The Western Journal of Medicine, v154, n1, p78(10)

Jan,

1991

PUBLICATION FORMAT: Magazine/Journal ISSN: 0093-0415 LANGUAGE: English

RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE: Professional

WORD COUNT: 8895 LINE COUNT: 00886

... goal at a lower cost. In selected patients with medication side effects, difficult-to-control **blood** pressure, or other undefined circumstances, however, **monitoring** may provide important information.

Clinical Trials for Mild Hypertension--Lessons

Lessons and Questions

Including the landmark Veterans Administration (VA) Cooperative Study

. . .

6/3,K/80 (Item 32 from file: 149)

DIALOG(R) File 149:TGG Health&Wellness DB(SM) (c) 2003 The Gale Group. All rts. reserv.

01254386 SUPPLIER NUMBER: 09301926 (USE FORMAT 7 OR 9 FOR FULL TEXT)
New uses and old for carbamazepine. (includes information on generics and on risk of birth defects)

Browne, Thomas R., III; Empting, Larry; Lydiard, R. Bruce; Yerby, Mark S. Patient Care, v24, n20, p48(15)

Dec 15,

1990

PUBLICATION FORMAT: Magazine/Journal ISSN: 0031-305X LANGUAGE: English

RECORD TYPE: Fulltext TARGET AUDIENCE: Professional

WORD COUNT: 6292 LINE COUNT: 00562

... that they are taking the drug). Dosage should be tapered when therapy is stopped.

The monitoring protocol used in many clinical trials of carbamazepine requires a complete blood count, liver function screen, and electrolyte determinations before starting treatment, then at the following intervals...

6/3,K/81 (Item 33 from file: 149)

DIALOG(R) File 149:TGG Health&Wellness DB(SM) (c) 2003 The Gale Group. All rts. reserv.

01206445 SUPPLIER NUMBER: 09265886

National high blood pressure education program working group report on ambulatory blood pressure monitoring.

Archives of Internal Medicine, v150, n11, p2265(11)

Nov,

1990

PUBLICATION FORMAT: Magazine/Journal ISSN: 0003-9926 LANGUAGE: English

RECORD TYPE: Abstract TARGET AUDIENCE: Professional

... ABSTRACT: those at higher risk of heart attack is being studied. It has

proved useful in research and clinical situations, such as monitoring the effects of drugs taken to lower high blood pressure, especially since blood pressure taken in the doctor's office may be affected by the timing of when...

(Item 34 from file: 149) 6/3,K/82 DIALOG(R) File 149:TGG Health&Wellness DB(SM) (c) 2003 The Gale Group. All rts. reserv.

(USE FORMAT 7 OR 9 FOR FULL TEXT) SUPPLIER NUMBER: 06526683 01147692 Brain imaging: applications in psychiatry.

Andreasen, Nancy C.

Science, v239, n4846, p1381(8)

March 18,

1988

ISSN: 0036-8075 LANGUAGE: English PUBLICATION FORMAT: Magazine/Journal

RECORD TYPE: Fulltext TARGET AUDIENCE: Academic

LINE COUNT: 00712 6846 WORD COUNT:

image construction.

Several 123I-labeled tracers, such as 123-I-IMP, are now available for clinical imaging trials in the United States. These tracers are ideal for labeling blood flow and may permit examination of perfusion in subcortical regions such as the basal ganglia...

(Item 1 from file: 444) 6/3,K/83 DIALOG(R) File 444: New England Journal of Med. (c) 2003 Mass. Med. Soc. All rts. reserv.

Copyright 1999 by the Massachusetts Medical Society

Oral versus Intravenous Empirical Antimicrobial Therapy for Fever in Patients with Granulocytopenia Who Are Receiving Cancer Chemotherapy (Original Articles)

Kern, Winfried V.; Cometta, Alain; de Bock, Robrecht; Langenaeken, John; Paesmans, Marianne; Gaya, Harold; Zanetti, Giorgio; Calandra, Thierry; Glauser, Michel P.; Crokaert, Francoise; Klastersky, Jean; Skoutelis, Athanasios; Bassaris, Harry; Zinner, Stephen H.; Viscoli, Claudio; Engelhard, Dan; Padmos, Andrew; for the International Antimicrobial Therapy Cooperative Group of the European Organization for Research and Treatment of Cancer. The New England Journal of Medicine Jul 29, 1999; 341 (5),pp 312-318 WORD COUNT: 05645 LINE COUNT: 00409

CITED REFERENCES

...patients with cancer. Antimicrob Agents Chemother 1996;40:1108-15.

- 28. Makuch R, Simon R. Sample size requirements for evaluating a conservative therapy. Cancer Treat Rep 1978;62:1037-40.
- trial progress. In: Pocock SJ. Clinical 29. Monitoring practical approach. Chichester, England: John Wiley, 1983:142-59. 30. O'Brien PC, Fleming...

(Item 2 from file: 444) 6/3,K/84 DIALOG(R) File 444: New England Journal of Med. (c) 2003 Mass. Med. Soc. All rts. reserv.

00102824

Copyright 1987 by the Massachusetts Medical Society

Of Oral Charcoal In Congenital Erythropoietic Therapeutic Efficacy Porphyria (Medical Intelligence)

Pimstone, Neville R.; Gandhi, Shobha N., M.B.B.S.; Mukerji, Sudhir K.,

The New England Journal of Medicine February 12, 1987; 316 (7),pp 390-393

WORD COUNT: 05193 LINE COUNT: 00376

...in consecutive clinical studies conducted two weeks apart. In both investigations, skin, urine, and plasma samples were obtained, and their porphyrin contents were monitored . In the first clinical patient ingested the activated charcoal as a slurry (30 g every 3 hours for...

(Item 3 from file: 444) 6/3,K/85 DIALOG(R) File 444: New England Journal of Med. (c) 2003 Mass. Med. Soc. All rts. reserv.

00100454

Copyright 1985 by the Massachusetts Medical Society

Prevention of Cardiac Disease by Subcutaneous Deferoxamine in Patients with Thalassemia Major (Original Article)

Wolfe, Lawrence; Olivieri, Nancy; Sallan, Darlene, R.N.; Colan, Steven; Rose, Vera; Propper, Richard; Freedman, Melvin H.; Nathan, David G. The New England Journal of Medicine June 20, 1985; 312 (25),pp 1600-1603 WORD COUNT: 03849 LINE COUNT: 00278

... All patients had previously undergone splenectomy. None had cardiac disease at the onset of the study, as judged by clinical examination, echocardiography, and Holter monitoring . On entrance into this study, patients began a program of supertransfusion (maintenance of blood hemoglobin levels no lower than 12 g per deciliter) and deferoxamine, 25 to 50 mg...

```
Set
        Items
                Description
· S1
      2223801
                TRACK? OR TRACE? OR TRACING OR MONITOR?
\mathbb{S}^2
      2451256
                 (CLINICAL? OR MEDICAL) (5N) (STUDIES OR STUDY OR RESEARCH? OR
              TRIAL?)
S3
     10360541
                SAMPLE? OR EXHIBIT? OR BLOOD? OR SERUM OR GENOTYP? OR PHEN-
             OTYP?
S4
       372354
                SCIENTIST? OR RESEARCHER? OR WORKER OR ATTENDANT?
S5
        13851
                S1(5N)S2
S6
         2947
                S5 (15N) S3
S7
            9
                S6(15N)S4
          292
S8
                S6(5N) (PROTOCOL? OR STEP? ? OR PROCEDUR?)
S9
          222
                S8 NOT PY>1999
S10
          208
                RD (unique items)
       237468
                CHECKLIST? OR (WORK OR CHECK) () LIST? ? OR ROSTER? ? OR ASS-
S11
             IGNMENT? OR WORKLIST?
S12
            0
                S10(20N)S11
                S10 AND S11
S13
            1
S14
            9
                S6 AND S11
S15
            9
                S13 OR S14
? show file
File
       2:INSPEC 1969-2003/Apr W2
         (c) 2003 Institution of Electrical Engineers
      35:Dissertation Abs Online 1861-2003/Mar
File
         (c) 2003 ProQuest Info&Learning
      65: Inside Conferences 1993-2003/Apr W3
File
         (c) 2003 BLDSC all rts. reserv.
      99:Wilson Appl. Sci & Tech Abs 1983-2003/Mar
File
         (c) 2003 The HW Wilson Co.
File 233: Internet & Personal Comp. Abs. 1981-2003/Mar
         (c) 2003 Info. Today Inc.
File 474: New York Times Abs 1969-2003/Apr 24
         (c) 2003 The New York Times
File 475: Wall Street Journal Abs 1973-2003/Apr 24
         (c) 2003 The New York Times
File 583:Gale Group Globalbase(TM) 1986-2002/Dec 13
         (c) 2002 The Gale Group
File 256:SoftBase:Reviews, Companies&Prods. 82-2003/Mar
         (c) 2003 Info. Sources Inc
File
       5:Biosis Previews(R) 1969-2003/Apr W3
         (c) 2003 BIOSIS
File
     73:EMBASE 1974-2003/Apr W3
         (c) 2003 Elsevier Science B.V.
File 155:MEDLINE(R) 1966-2003/Apr W3
         (c) format only 2003 The Dialog Corp.
     34:SciSearch(R) Cited Ref Sci 1990-2003/Apr W3
         (c) 2003 Inst for Sci Info
File 434:SciSearch(R) Cited Ref Sci 1974-1989/Dec
```

(c) 1998 Inst for Sci Info

(Item 1 from file: 5) DIALOG(R) File 5:Biosis Previews(R) (c) 2003 BIOSIS. All rts. reserv. BIOSIS NO.: 000093116934 08117586 INTERIM ANALYSES FOR MONITORING CLINICAL TRIALS THAT DO NOT MATERIALLY AFFECT THE TYPE I ERROR RATE AUTHOR: GOULD A L AUTHOR ADDRESS: MERCK SHARP AND DOHME RES. LAB., WEST POINT, PA. 19486, JOURNAL: STAT MED 11 (1). 1992. 55-66. 1992 FULL JOURNAL NAME: Statistics in Medicine CODEN: SMEDD RECORD TYPE: Abstract LANGUAGE: ENGLISH trials often requires examining the ABSTRACT: Monitoring clinical interim findings to see if the sample size originally specified in the protocol will provide the required power against the null hypothesis when the alternative hypothesis is true, and to increase the sample size if necessary. This paper presents a new method, based on the overall response rate, for carrying out interim power evaluations when the observations have binomial distributions, without unblinding the treatment assignments of materially affecting the type I error rate. Simulation study results confirm the performance of the method. DESCRIPTORS: HUMAN TREATMENT CONCEPT CODES: Public Health-Public Health Administration and Statistics 37010 Public Health-Health Services and Medical Care 37012 04500 Mathematical Biology and Statistical Methods Pathology, General and Miscellaneous-Therapy (1971-) 12512 BIOSYSTEMATIC CODES: Hominidae 86215 BIOSYSTEMATIC CLASSIFICATION (SUPER TAXA): Animals Chordates Vertebrates Mammals Primates Humans (Item 1 from file: 73) 15/5/2 DIALOG(R) File 73: EMBASE (c) 2003 Elsevier Science B.V. All rts. reserv. EMBASE No: 2002406578 11830278 Home sodium monitoring in patients with diabetes insipidus Green R.P.; Landt M. Dr. R.P. Green, One Children's Place, Box 8116, St Louis, MO 63110 United States Journal of Pediatrics (J. PEDIATR.) (United States) 01 NOV 2002, 141/5 (618-624) CODEN: JOPDA ISSN: 0022-3476 DOCUMENT TYPE: Journal ; Article SUMMARY LANGUAGE: ENGLISH LANGUAGE: ENGLISH

NUMBER OF REFERENCES: 15

Objective: To determine whether home care givers can accurately measure plasma sodium in children with diabetes insipidus (DI) by using an I-STAT portable clinical analyzer (PCA) and to collect preliminary data on home PCA use. Study design: Care givers of 4 children with DI and impaired thirst or inability to access water freely were instructed in PCA use. During an initial preclinical phase, the accuracy of sodium concentration measured by care givers was assessed by comparison to simultaneous analysis in a clinical laboratory. Participants were subsequently randomly assigned to daily home PCA monitoring or routine care. All participants crossed over from their original randomized group assignment to the alternate group. Results: After a single education session, all care givers were able to perform PCA testing. There was good correlation between PCA and laboratory sodium (r = 0.92). On the basis of Error Grid Analysis, use of the PCA sodium would have resulted in treatment decisions identical to those made based on the laboratory sodium value in 62 of 66 instances. Four minor differences in treatment would have occurred. There was no statistically significant difference in clinical outcome during daily monitoring versus routine care. Conclusions: Results obtained by care givers using the PCA are sufficiently reliable for assessment of fluid status and making treatment decisions.

DEVICE BRAND NAME/MANUFACTURER NAME: Portable Clinical Analyzer/i-STAT/ United States DEVICE MANUFACTURER NAMES: i-STAT/United States DRUG DESCRIPTORS: *sodium--endogenous compound--ec MEDICAL DESCRIPTORS: *diabetes insipidus sodium blood level; caregiver; home monitoring; accuracy; device infection; clinical laboratory; randomization; decision making; outcomes research; reliability; human; male; female; case report; clinical trial; crossover procedure; child; article; priority journal CAS REGISTRY NO.: 7440-23-5 (sodium) SECTION HEADINGS: 007 Pediatrics and Pediatric Surgery

027 Biophysics, Bioengineering and Medical Instrumentation

(Item 2 from file: 73) 15/5/3 DIALOG(R)File 73:EMBASE (c) 2003 Elsevier Science B.V. All rts. reserv.

EMBASE No: 1998173464

A comparison of intravascular pressure monitoring system contamination and patient bacteremia with use of 48- and 72-hour system change intervals McLane C.; Morris L.; Holm K.

C. McLane, CCRN, 6240 South Rutherford, Chicago, IL 60638 United States Heart and Lung: Journal of Acute and Critical Care (HEART LUNG J. ACUTE CRIT. CARE) (United States) 1998, 27/3 (200-208)

ISSN: 0147-9563 CODEN: HELUA DOCUMENT TYPE: Journal; Article

SUMMARY LANGUAGE: ENGLISH LANGUAGE: ENGLISH

NUMBER OF REFERENCES: 23

OBJECTIVE: To determine the incidence of culture positivity in intravascular monitoring systems by comparing 48- versus 72-hour intervals for flush solution, stopcocks, and catheters on removal. DESIGN: Prospective, quasi-experimental, random assignment . SETTING: Intensive care units of a midwestern university medical center and a community hospital. PATIENTS: Seventy-six critically ill adult patients, ranging in age from 24 to 96 years (X = 61.6), requiring arterial or pulmonary artery catheters. OUTCOME MEASURE: Culture positivity of flush solution, stopcocks, or catheter tips. INTERVENTION: Data collection was initiated at designated change intervals of 48- or 72-hours; cultures were taken of flush solution and stopcocks; catheter tip cultures were obtained on catheter removal. RESULTS: Chi-square analyses indicated that increasing the change interval to 72 hours resulted in no significant difference in culture positivity of catheter tips. However, the difference between the 48- and 72-hour groups in culture-positivity rates of stopcocks from arterial catheters was significant (1, N = 82) = 6.86, p less than 0.01. CONCLUSIONS: Our results showed that increasing the change interval to 72 hours did not increase the risk of catheter-associated infection or catheter-associated bacteremia. Chi-square analysis did not show an association between culture-positive stopcocks, the incidence of culturepositive catheter tips, entries into the system, or catheter-related bacteremia and a change interval that was increased to 72 hours. Thus, increasing the change interval to 72 hours does not increase the risk of infection.

MEDICAL DESCRIPTORS:

*bacteremia--complication--co; *bacteremia--diagnosis--di; *monitor; *
infection--complication--co; *infection--diagnosis--di
bacterium culture; incidence; artery catheter; infection risk; bacterium
contamination; blood pressure monitoring; human; male; female; major
clinical study; clinical trial; randomized controlled trial;
controlled study; aged; adult; article; priority journal
SECTION HEADINGS:

- 004 Microbiology: Bacteriology, Mycology, Parasitology and Virology
- 006 Internal Medicine
- 027 Biophysics, Bioengineering and Medical Instrumentation
- 037 Drug Literature Index

15/5/4 (Item 3 from file: 73)

DIALOG(R)File 73:EMBASE

(c) 2003 Elsevier Science B.V. All rts. reserv.

06918080 EMBASE No: 1997202542

Trial of calcium to prevent preeclampsia

Levine R.J.; Hauth J.C.; Curet L.B.; Sibai B.M.; Catalano P.M.; Morris C.D.; Dersimonian R.; Esterlitz J.R.; Raymond E.G.; Bild D.E.; Clemens J.D.; Cutler J.A.

Dr. R.J. Levine, DESPR, NICHHD, National Institutes of Health, Bethesda, MD 20892-7510 United States

New England Journal of Medicine (NEW ENGL. J. MED.) (United States) 1997, 337/2 (69-76)

CODEN: NEJMA ISSN: 0028-4793 DOCUMENT TYPE: Journal; Article

LANGUAGE: ENGLISH SUMMARY LANGUAGE: ENGLISH

NUMBER OF REFERENCES: 36

Background Previous trials have suggested that calcium supplementation during pregnancy may reduce the risk of preeclampsia. However, differences in study design and a low dietary calcium intake in the populations studied limit acceptance of the data. Methods. We randomly assigned 4589 healthy nulliparous women who were 13 to 21 weeks pregnant to receive daily treatment with either 2 g of elemental calcium or placebo for the remainder of their pregnancies. Surveillance for preeclampsia was conducted by personnel unaware of treatment-group assignments, using standardized measurements of blood pressure and urinary protein excretion at uniformly scheduled prenatal visits, protocols for monitoring these measurements during the hospitalization for delivery, and reviews of medical records of

unscheduled outpatient visits and all hospitalizations. Results Calcium supplementation did not significantly reduce the incidence or severity of preeclampsia or delay its onset. Preeclampsia occurred in 158 of the 2295 women in the calcium group (6.9 percent) add 168 of the 2294 women in the placebo group (7.3 percent) (relative risk, 0.94; 95 percent confidence interval, 0.76 to 1.16). There were no significant differences between the two groups in the prevalence of pregnancy-associated hypertension without preeclampsia (15.3 percent vs. 17.3 percent) or of all hypertensive disorders (22.2 percent vs. 24.6 percent). The mean systolic and diastolic blood pressures during pregnancy were similar in both groups. Calcium did not reduce the numbers of preterm deliveries, small-for-gestational-age births, or fetal and neonatal deaths; nor did it increase urolithiasis during pregnancy. Conclusions Calcium supplementation during pregnancy did not prevent preeclampsia, pregnancy-associated hypertension, or adverse perinatal outcomes in healthy nulliparous women.

DRUG DESCRIPTORS:
*calcium--clinical trial--ct; *calcium--drug therapy--dt
MEDICAL DESCRIPTORS:
*calcium intake; *preeclampsia--prevention--pc; *preeclampsia--drug therapy
--dt
adult; article; blood pressure monitoring; clinical trial;
controlled study; diet supplementation; disease severity; female; follow
up; human; major clinical study; oral drug administration; prenatal care;
priority journal; protein urine level; randomized controlled trial;
treatment outcome
CAS REGISTRY NO.: 7440-70-2 (calcium)
SECTION HEADINGS:
010 Obstetrics and Gynecology
037 Drug Literature Index

15/5/5 (Item 4 from file: 73)
DIALOG(R)File 73:EMBASE
(c) 2003 Elsevier Science B.V. All rts. reserv.

06352968 EMBASE No: 1996017122

United Kingdom prospective diabetes study 17: A 9-year update of a randomized, controlled trial on the effect of improved metabolic control on complications in non-insulin-dependent diabetes mellitus

Turner R.; Cull C.; Holman R.

United Kingdom PDSG, Diabetes Research Laboratories, Radcliffe Infirmary, Woodstock Road, Oxford OX2 6HE United Kingdom

Annals of Internal Medicine (ANN. INTERN. MED.) (United States) 1996,

124/1 II (136-145) CODEN: AIMEA ISSN: 0003-4819

DOCUMENT TYPE: Journal; Conference Paper

LANGUAGE: ENGLISH SUMMARY LANGUAGE: ENGLISH

Purpose: To report the progress (after 9-year follow-up) of a study designed to determine whether improved glucose control in patients with newly diagnosed non-insulin-dependent diabetes mellitus (NIDDM) is effective in reducing the incidence of clinical complications. Data Source: A multicenter, randomized, controlled trial of different therapies for NIDDM. After initial diet therapy, 4209 asymptomatic patients who remained hyperglycemic (fasting plasma glucose levels, 6.0 to 15.0 mmol/L) were assigned to either a conventional therapy policy, primarily with diet alone, or to an intensive therapy policy, aiming for fasting plasma glucose levels of less than 6.0 mmol/L, with assignment to primary therapy with sulfonylurea or insulin (which increased insulin supply) or metformin (which enhanced insulin sensitivity). Results: All three modes of

pharmacologic therapy in the intensively treated group-sulfonylurea, insulin, and metformin-had similar efficacy in reducing the fasting plasma glucose and glycated hemoglobin levels. Over 9 years, patients assigned to intensive therapy with sulfonylurea or insulin had lower fasting plasma glucose levels (median, 7.3 and 9.0 mmol/L, respectively) and lower hemoglobin A(1c) levels (6.7% and 7.5%, respectively) than patients assigned to conventional therapy. Regardless of the assigned therapy, however, the fasting plasma glucose and hemoglobin A(1c) levels increased, and maintaining near-normal glycemia was, in general, not feasible. Even insulin therapy did not achieve the therapeutic goal of near-normal glycemia because of the difficulty in treating marked hyperglycemia and the risk for hypoglycemic episodes. Nine years after the diagnosis of diabetes, 29% of the patients had had a diabetes-related clinical end point, 20% had had a macrovascular complication, and 9% had had a microvascular complication. Conclusions: A report will be published in 1998 after a median duration from randomization of 11 years (range, 6 to 20 years) with an 81% power at a 1% level of significance of detecting whether the obtained improvement in glucose control causes a 15% decrease or increase in the incidence of major complications and whether any specific therapy is advantageous or disadvantageous.

```
DRUG DESCRIPTORS:
*chlorpropamide--clinical trial--ct; *chlorpropamide--drug comparison--cm;
*chlorpropamide--drug therapy--dt; *glibenclamide--clinical trial--ct; *
glibenclamide--drug therapy--dt; *glibenclamide--drug comparison--cm; *
glipizide--clinical trial--ct; *glipizide--drug comparison--cm; *glipizide
--drug therapy--dt; *insulin zinc suspension--drug therapy--dt; *insulin
zinc suspension--drug dose--do; *insulin zinc suspension--drug combination
--cb; *insulin zinc suspension--clinical trial--ct; *insulin zinc
suspension--drug comparison--cm; *isophane insulin--clinical trial--ct; *
isophane insulin--drug combination--cb; *isophane insulin--drug comparison
--cm; *isophane insulin--drug dose--do; *isophane insulin--drug therapy--dt
; *metformin--clinical trial--ct; *metformin--drug combination--cb; *
metformin--drug therapy--dt
atenolol--drug therapy--dt; captopril--drug therapy--dt
MEDICAL DESCRIPTORS:
*diabetic angiopathy--complication--co; *diabetic angiopathy--prevention
--pc; *diabetic nephropathy--prevention--pc; *diabetic nephropathy
--complication--co; *diabetic retinopathy--prevention--pc; *diabetic
retinopathy--complication--co; *non insulin dependent diabetes mellitus
--epidemiology--ep; *non insulin dependent diabetes mellitus--drug therapy
--dt; *non insulin dependent diabetes mellitus--therapy--th; *non insulin
dependent diabetes mellitus--etiology--et
adult; blood glucose monitoring; clinical
                                                 trial; conference paper;
controlled study; diabetic diet; dose response; drug efficacy; drug
potency; female; hemoglobin determination; human; hypertension
--complication--co; hypertension--drug therapy--dt; insulin treatment;
major clinical study; male; multicenter study; priority journal; randomized
controlled trial; treatment outcome; treatment planning
CAS REGISTRY NO.: 94-20-2 (chlorpropamide); 10238-21-8 (glibenclamide);
    29094-61-9 (glipizide); 8049-62-5 (insulin zinc suspension); 9004-17-5
    (isophane insulin); 1115-70-4, 657-24-9 (metformin); 29122-68-7 (
    atenolol); 62571-86-2 (captopril)
SECTION HEADINGS:
  003 Endocrinology
  012 Ophthalmology
  018 Cardiovascular Diseases and Cardiovascular Surgery
  028 Urology and Nephrology
  037 Drug Literature Index
```

```
(Item 5 from file: 73)
DIALOG(R) File 73: EMBASE
(c) 2003 Elsevier Science B.V. All rts. reserv.
            EMBASE No: 1995120871
 Parent-rated family climate: A concomitant to metabolic control in
juvenile IDDM?
  Hansson K.; Ryden O.; Johnsson P.
  Dept. of Child and Youth Psychiatry, University of Lund, P.O. Box
  638, S-22009 Lund Sweden
  Family Systems Medicine ( FAM. SYST. MED. ) (United States) 1994, 12/4
  (405 - 413)
                ISSN: 0736-1718
 CODEN: FSMEE
  DOCUMENT TYPE: Journal; Article
  LANGUAGE: ENGLISH SUMMARY LANGUAGE: ENGLISH
  Parents in 25 families with a diabetic child in poor metabolic control
rated their family climate with a checklist containing 85 adjectives
constituting four climate factors: Closeness, Distance, Expressiveness, and
Chaos. Data were obtained before and after one of three treatment
conditions: family therapy, pediatric support, and conventional treatment.
Data also were obtained from 12 patients in optimal metabolic control and
their families, and from 10 reference families with healthy children. Prior
to treatment, the parents of children in poor metabolic control rated their
family climate as lower on Closeness and higher on Chaos, as compared with
the parents of children in optimal control. No differential effects on
family climate were found between the family therapy and pediatric support
groups. The combined treatment groups rated themselves as higher on
Closeness and lower on Chaos after intervention than before. We concluded
that parents with children in poor metabolic control normalize their
self-treated family climate following family treatment.
DRUG DESCRIPTORS:
*hemoglobin alc--endogenous compound--ec
MEDICAL DESCRIPTORS:
*family life; *family therapy; *insulin dependent diabetes mellitus; *
juvenile diabetes mellitus
adolescent; adult; article; blood glucose monitoring; child; child
parent relation; clinical article; clinical trial; conflict;
controlled study; diabetes control; emotion; female; human; intimacy;
male; randomized controlled trial; self concept
CAS REGISTRY NO.: 62572-11-6 (hemoglobin alc)
SECTION HEADINGS:
  003 Endocrinology
  006 Internal Medicine
  007 Pediatrics and Pediatric Surgery
  017 Public Health, Social Medical and Epidemiology
  032 Psychiatry
            (Item 6 from file: 73)
 15/5/7
DIALOG(R)File 73:EMBASE
(c) 2003 Elsevier Science B.V. All rts. reserv.
             EMBASE No: 1992198665
05058449
  Nonpharmacologic intervention to reduce blood pressure in older patients
with mild hypertension
  Applegate W.B.; Miller S.T.; Elam J.T.; Cushman W.C.; El Derwi D.; Brewer
```

Bode Akintola 25-Apr-03

66 N Pauline, Memphis, TN 38105 United States

A.; Granev M.J.

Archives of Internal Medicine (ARCH. INTERN. MED.) (United States)

1992, 152/6 (1162-1166)

CODEN: AIMDA ISSN: 0003-9926 DOCUMENT TYPE: Journal; Article

LANGUAGE: ENGLISH SUMMARY LANGUAGE: ENGLISH

Background. - Although nonpharmacologic interventions are widely recommended in the therapy of high blood pressure in older adults, surprisingly little data exist to confirm the efficacy of these interventions in older persons. Methods. - We conducted a randomized, controlled clinical trial in persons aged 60 to 85 years with a diastolic blood pressure of 85 to 100 mm Hg. The experimental arm was a nonpharmacologic intervention combining weight reduction, sodium restriction, and increased physical activity. The nonpharmacologic intervention consisted of eight weekly group and two individual sessions during the intensive phase, followed by four monthly group sessions during the maintenance phase. The control group received no treatment during the study. Blood pressure was assessed by certified technicians (blinded to group assignment) using random zero sphygmomanometers. Results. - Of 56 participants randomized, 47 completed the entire 6-month trial (21 in the intervention group and 26 in the control group). Attendance at the intervention sessions was excellent. The intervention group lost more weight (-2.1 kg) over 6 months than the control group (+0.3 kg). Trends for decreasing 24-hour urine sodium excretion in both the intervention and control groups, with greater trend in the intervention group, were not statistically significant. The intervention group experienced more reduction in systolic and diastolic blood pressure than did the control group (mean differences between groups at 6 months, 4.2/4.9 mm Hg, respectively). Conclusions. - Our data indicate that a nonpharmacologic intervention will lower systolic and diastolic blood pressure levels in older people with borderline or mild elevations of diastolic blood pressure.

MEDICAL DESCRIPTORS:

*aging; *blood pressure regulation; *hypertension--therapy--th adult; aged; article; blood pressure monitoring; clinical article; controlled study; diastolic blood pressure; female; human; male; physical activity; priority journal; sodium restriction; sodium urine level; weight reduction SECTION HEADINGS:

006 Internal Medicine

017 Public Health, Social Medical and Epidemiology

15/5/8 (Item 1 from file: 155)

DIALOG(R) File 155: MEDLINE(R)

(c) format only 2003 The Dialog Corp. All rts. reserv.

07350149 92213308 PMID: 1557576

Interim analyses for monitoring clinical trials that do not materially affect the type I error rate.

Gould A L

Merck, Sharp, and Dohme Research Laboratories, West Point, PA 19486. Statistics in medicine (ENGLAND) Jan 15 1992, 11 (1) p55-66, ISSN 0277-6715 Journal Code: 8215016

Document type: Journal Article

Languages: ENGLISH
Main Citation Owner: NLM
Record type: Completed
Subfile: INDEX MEDICUS

Monitoring clinical trials often requires examining the interim

Bode Akintola 25-Apr-03

findings to see if the **sample** size originally specified in the protocol will provide the required power against the null hypothesis when the alternative hypothesis is true, and to increase the sample size if necessary. This paper presents a new method, based on the overall response rate, for carrying out interim power evaluations when the observations have binomial distributions, without unblinding the treatment **assignments** or materially affecting the type I error rate. Simulation study results confirm the performance of the method.

Tags: Human

Descriptors: *Clinical Trials--statistics and numerical data--SN; Bias (Epidemiology); Data Interpretation, Statistical; Models, Statistical; Odds Ratio

Record Date Created: 19920504
Record Date Completed: 19920504

15/5/9 (Item 1 from file: 34)

DIALOG(R)File 34:SciSearch(R) Cited Ref Sci (c) 2003 Inst for Sci Info. All rts. reserv.

01488332 Genuine Article#: HC732 Number of References: 0

(NO REFS KEYED)

Title: INTERIM ANALYSES FOR MONITORING CLINICAL-TRIALS THAT DO NOT MATERIALLY AFFECT THE TYPE-I ERROR RATE

Author(s): GOULD AL

Corporate Source: MERCK SHARP & DOHME LTD/W POINT//PA/19486 Journal: STATISTICS IN MEDICINE, 1992, V11, N1 (JAN 15), P55-66

Language: ENGLISH Document Type: ARTICLE

Geographic Location: USA

Subfile: SciSearch; CC LIFE--Current Contents, Life Sciences; CC CLIN--Current Contents, Clinical Medicine

Journal Subject Category: MEDICINE, RESEARCH & EXPERIMENTAL; MATHEMATICS, APPLIED; PUBLIC HEALTH

Abstract: Monitoring clinical trials often requires examining the interim findings to see if the sample size originally specified in the protocol will provide the required power against the null hypothesis when the alternative hypothesis is true, and to increase the sample size if necessary. This paper presents a new method, based on the overall response rate, for carrying out interim power evaluations when the observations have binomial distributions, without unblinding the treatment assignments or materially affecting the type I error rate. Simulation study results confirm the performance of the method.